

**PRELIMINARY CATALOG OF DESCRIBED
SPECIES OF THE GENUS NERITA
(GASTROPODA: NERITIDAE)**

By

Lucius G. Eldredge

University of Guam Marine Laboratory

Technical Report No. 89

August 1987

**PRELIMINARY CATALOG OF DESCRIBED SPECIES OF THE
GENUS NERITA (GASTROPODA: NERITIDAE)**

By

Lucius G. Eldredge

University of Guam Marine Laboratory

Technical Report No. 89

August 1987

ACKNOWLEDGEMENTS

This catalog could not have materialized without the full cooperation and assistance of a wide variety of places and people. A number of private and public libraries have been most kind. Two libraries at the Smithsonian Institution in Washington, D.C. have been particularly helpful. These are the general institutional Library and the Department of Mollusks library. The Mollusk Library at the Museum of Comparative Zoology, Harvard University, allowed me to spend two days copying pertinent and otherwise unavailable literature. Libraries of both the University of Washington (Seattle) and the University of Maryland (College Park) have been searched for information. In Hawaii, literature has been acquired at both the B. P. Bishop Museum and University of Hawaii libraries; the University of Guam library has also been utilized. Personally, Dr. G. J. Vermeij, University of Maryland has made substantial contributions and greatly assisted in this study. Additional thanks are extended to Jo Eldredge who managed to remain calm during the compilation and final editing of this catalog.

This report is a thesis submitted to the Graduate School of the University of Hawaii in partial fulfillment of the requirements for the M.L.S. degree

ABSTRACT

Through a review of the major nomenclatorial works, a total of 780 species names were recorded, all having been described within the genus Nerita of the archaeogastropod family Neritidae. Some 279 were originally described from fossil specimens; the remaining are Recent. This catalog includes the full name and full bibliographic citation, as originally published. The collection location of the original specimen(s) is provided when known, and remarks are presented when appropriate. More than 100 of the Recent names are invalid, being nomen nudum, nomen dubium, or rejected.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	iii
ABSTRACT	iv
LIST OF TABLES	vi
I. INTRODUCTION	1
II. MATERIALS AND METHODS	2
III. RESULTS, DISCUSSION AND CONCLUSIONS	6
LITERATURE CITED	159

LIST OF TABLES

Table	Page
1 Catalog of Recent <u>Nerita</u> Species	8
2 List of Fossil <u>Nerita</u> Species	126
3 <u>Nerita</u> Species Names Described More Than Once	156

I. INTRODUCTION

The archaeogastropod family Neritidae is composed of several genera. Representatives of the family are a major component of the coral reef ecosystems of the world. Most species live in the high intertidal zone; few remain submerged for long periods of time. In its entirety, this is only molluscan family with species found in marine, freshwater, brackish, and terrestrial habitats.

The genus Nerita contains the greatest number of described species and has the longest taxonomic history in this family, having originally been described by Linnaeus in 1758. This genus was reviewed and updated by Hanley (1855). Since its original description, more than 750 species have been proposed as belonging to Nerita. However, many of the names are currently either invalid or rejected for a variety of different reasons through the implementation of the International Code of Zoological Nomenclature (International Commission for Zoological Nomenclature, 3rd ed., 1985).

The objectives of this study have been to document all of the described species as thoroughly as possible and to catalog them in a single, consistent manner. Additionally, the type locality and the location of the type specimen(s) have been noted where known. Remarks have been made as to the present status of each species, since many species names belong to individuals which are clearly not members of the genus Nerita.

II. MATERIALS AND METHODS

In zoology the definitive work concerning the validity of a specific name is the original description of that species. Since the work of Linnaeus (1758), names have been binomial; that is, each species has been described under a combination of two names, the first being the genus and the second, the species itself. For nearly a century and a half, however, no consistent standard governed the validity of the names of new species described. Begun in 1905, as International Rules of Zoological Nomenclature, the present International Code of Zoological Nomenclature (International Commission for Zoological Nomenclature, 3rd ed., 1985) now stipulates criteria for the myriad of details affecting recognition of names as valid and the citability of their publication.

To compile a catalog of described species of any group, the original descriptions must be consulted, for it is upon these descriptions that specific distinctions are based. Merely locating the citations can be problematic for some groups, as not many direct methods exist through which these citations can be obtained. Sherborn's (1902-1933) two-series, multivolume Index Animalium Sive Index Nomonim Quae ab A.D. MDCCLVIII Generibus et Speciebus Animalium Imposita Sunt covers all animal species described between 1758, the starting date for binomial nomenclature, and 1850. The monumental index series Zoological Record has covered all

zoological literature published since 1864 and continues to the present day. Thus, there was an approximate 14-year gap in the compilation of species descriptions. The Index to the Species of Mollusca Introduced from 1850 to 1870 was compiled by Ruhoff (1980) under the auspices of the Smithsonian Institution to cover this gap for the mollusks.

From among these four works a basic listing was made of all mollusk citations between 1758 and the present. From this list a single compilation of descriptions of Nerita species has been made. A search for the original material was initiated through visits to several U.S. libraries, and the majority of the original descriptions has been accumulated. Most of the original material was initially photocopied so that it could be used at a later time.

The format of the catalog is designed to include as much information as possible. The species are listed in alphabetical order. Following each, a full bibliographic citation is given in the format of the CBE Style Manuel (Council of Biology Editors, 1972). The original description as published has been quoted exactly. The majority of these descriptions are in Latin and have not been translated for this catalog. Apparent style differences herein reflect differences in the exact wording of the original descriptions. An example is *Nerita*, *Ner.*, or *N.*, pertaining to the genus within which the species is located.

—

The type locality follows. This is the collection location of the original specimen(s). In most of the older literature the exact locality or habitat is either not given or was not known. In many, expressions such as "Insulae Americae" or "O. Asiae" are stated. These "geographic locations" are of course useless to the modern zoologist who would like to look at the area from which specimens might have been collected.

Remarks follow, as appropriate. Notes on current validity are included where known, which, according to the Code, may be:

Nomen dubium: A descriptive term meaning name of unknown or doubtful application (p. 260).

Nomen nudum: A name that, if published before 1931, fails to conform to Article 12; or, if published after 1930, fails to conform to Article 13. A nomen nudum is not an available name and therefore the same name may be made available later for the same or a different concept; in such a case it would take authorship and date from that act of establishment, not from any earlier publication as nomen nudum (p. 260).

Rejected: Any name, available, unavailable, or vernacular, for a taxon other than its valid name [following the Official Index of Rejected and Invalid Works in Zoological Nomenclature (Hemming, 1958)] (p. 259).

Where known, the location of the type specimen(s) is given. The type specimen(s) is the actual individual(s) upon which the original description is based. Type specimens must now be officially designated under the options of

the Code. Most recent descriptions contain this information; most older publications do not. Unfortunately, type specimens were not designated in most of the nineteenth century literature.

The catalog includes only those species which are thought to be of Recent geologic origin; that is, they are species which can be found living at the present time. Geologically older specimens are considered to be fossils. A list of fossil species has also been compiled. This list contains only the bibliographic citation for the original description, which is often incomplete, for the majority has not been examined. These are marked with an asterisk (*). The fossil species do not enter into the contemporary environment, and many of the descriptions are incomplete because of the nature of the available fossil material.

III. RESULTS, DISCUSSION AND CONCLUSIONS

A total of 780 species names have been recorded. Of these, 44 original descriptions could not be located, and those names are noted with an asterisk (*) at the citation.

Table 1 contains the full catalog of species of Recent, nonfossil Nerita. The full scientific name and full bibliographic citation are given in this Table. The description is copied as it originally appeared, and the location of the collection of specimens is listed where known. The remarks provided include additional information about the species or the species name. In an analysis of some of the names provided, 100 are considered invalid as either nomen nudum or nomen dubium. One work with four specific names has been officially rejected.

Table 2 contains a list of 279 species of Nerita which were described as fossils. Fossil records are sketchy and the resulting names are problematic.

Table 3 lists 36 species names which have been described more than once, thus, making subsequent names invalid in most cases.

The genus Nerita has a long history. Originally, all similar morphological types were placed in this genus. Linnaeus (1758) described 24 such species. Interestingly, at least ten of those names are still considered valid. Gmelin (1791) described 41 new species and listed a total of 72 names, of which 22 were those of Linnaeus. The most

prolific writer of the mid-nineteenth century was Recluz, who described 118 new species, many of which are still considered valid.

This catalog represents the only compilation of this nature for Nerita and one of very few for any of the gastropods. Future studies will include the location and the actual examination of the type specimens, leading to a total review and revision of the genus.

TABLE 1
CATALOG OF RECENT NERITA SPECIES

Nerita achatina

Reeve, L. A. 1855. *Conchologia iconica: or, illustrations of the shells of molluscous animals* 9 (*Nerita*), plate 16, species 68.

Description: *Ner. testa ovata, spira subexerserta, anfractibus angustis, convexis, versus aperturam latioribus, impresse linearis-sulcatis, sulcis valde irregularibus, labro intus lirato, columella declivi, obsolete granulata, dentata; carbonario-nigra, nitente, sulcis linearibus pallidis.*

Type locality: Not given.

Nerita aculeata

Gmelin, J. F. 1791. *Systema naturae*, 13th edition, p. 3686.

Description: *N. testa nigrescente transversim striata: striis spinosis, labio planiusculo laevi subdentato.*

Type locality: *Indiae fluviis, puligerae affinis.*

Remarks: See Mienis (1973); Neritina juttingae new name.

Nerita aculeata

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 193.

Description: *Nerita testa turrita, fusca, tuberculosa; anfractibus muricatis; labro depresso, crenulato.*

Type locality: *In paludosis Africæ torridæ.*

Nerita acupictus

Martyn, T. 1789. *Universal conchyologist*, Vol. 3, p. 107.

Remarks: Nonbinomial, invalid name; need to check "Rejected Works."

Nerita adansoniana

Recluz, M. C. A. 1841. *Description de quelques nouvelles espèces de Nerites vivantes. Rev. Zool. Soc. Cuvier.* 1841, p. 313.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. adansoniana, cont.

Description: Testa ovato-obtusa, laevi sub cuticula rufescenti albida, lineis, liturisque nigris, undulatis, angulosis aut intextis picta fasciataque; spira priminula convexo-rotundatal apice erosa; apertura subobliqua; labio convexiusculo, calloso, medio subarcuato et obsolete crenato.

Type locality: Le confluent du fleuve Senegal.

Nerita adenensis

Menis, H. K. 1978. Notes on recent and fossil Neritidae
8. Nerita adenensis, a new species from the Arabian Gulf.
Argamon, Israel J. Malacol. 6(3-4), p. 32.

Description [Diagnosis]: A multicolored, medium-sized Nerita, characterized by a completely smooth and highly polished operculum.

Type locality: Crater Beach, Aden.

Location of type: Holotype and 63 paratypes--Rijksmuseum van Natuurlijke Historie, Leiden, Holland.

Nerita adspersa

Recluz, M. C. A. 1853. Description de quelques nerites et d'une natice. J. Conchyliol. 4, p. 319.

Description: N. testa ovata, obliquata, rugosiuscula, nigricante, punctis albo-lutescentibus adspersa; anfractibus tribus supremis spiram depresso efformatis; apertura subquadrata, albida; labio angusto, valde declivi, crenulato, in medio arcuato et dente valido instructo.

Type locality: Unknown.

Nerita affinis

Gmelin, J. F. 1791. Systema naturae, 13th edition,
p. 3675.

Description: N. testa globosa crassiore: spira submucronata; anstractibus tribus.

Type locality: Oceano septentrionali.

Remarks: See Cernohorsky, 1974.

TABLE 1.. (Continued) CATALOG OF RECENT NERITA SPECIESNerita affinis

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 18, species 81.

Description: Not given.

Type locality: Not given.

Remarks: Species figured but not described in Reeve; Vermeij--synonym of N. guamensis.

Nerita africana

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 198.

Description: Ner. testa ovato-conoidea, subepidermide nigrescente strigis nigris longitudinalibus creberimmiss et maculis oblongis vel ovatis transversis, luteis, antice acutis et nigro marginatis, subregulatibus, undique picta; anfractibus tribus subconicus; infimo superne rugis raris notato; spira vix prominula, apice erosa; apertura obliqua, extus semi-oblonga, intus albido-cinerascente; labio convexiusculo superne calloao, antice plano rectiusculo et laevissime crenato.

Type locality: "Island of Fernando Po, west coast of Africa; found in a small stream by Capt. Downs, R.N."

Remarks: Described in the subgenus Neritina.

Nerita alabastrina

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

Nerita ala-papilionis

*Salis, C. U. von. 1793. [not seen]

Nerita alba

Adams, J. 1797. The specific characters of some minute shells discovered on the coast of Pembrokshire, with an account of a new animal. Trans. Linnaean Soc. 3, p. 67.

Description: N. testa laevi subpellucida, duobus anfractibus.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

N. alba, cont.

Type locality: Not given.

Nerita albicilla

Linnaeus, C. 1758. *Systema naturae*, p. 778.

Description: *N. testa striata, labiis subdentatis: interiore tuberculato.*

Type locality: Hitoe [Mollucas]

Nerita albipunctata

Reeve, L. A. 1855. *Conchologia iconica: or, illustrations of the shells of molluscous animals* 9 (*Nerita*), plate 13, species 61.

Description: *Ner. testa, ovato-semiglobosa, spira sub-exserta, obtusa, anfractibus angustis, superne concavo-depressis, spiraliter liratis, liris rude crenulatis, interstitiis punctatis, columella plano-excavata, sparsim granulata, obsolete minutissime denticulata; coeruleo-nigra, lirarum interstitiis albipunctatis.*

Type locality: Not given.

Nerita albolineata

Bolten, J. F. 1798. *Museum Boltenianum, Part 2*, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita albumen

Linnaeus, C. 1758. *Systema naturae*, p. 776.

Description: *N. testa umbilicata convexa, umbilico subcordato: lobo explanato.*

Type locality: East Asia [O. Asiae].

Remarks: Vermeij--now Polinices (Neverita) albumen.

Nerita alticola

Pilsbry, H. A. 1932. A Jamaican fluviatile Nerita.
Proc. Acad. Nat. Sci. Philadelphia 84, p. 11.

Description: The shell is globose, with very short, low

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. alticola, cont.

spire, of nearly 3 1/2 whorls, the last descending more rapidly in the last half turn and narrowly flattened below the suture. Surface smooth except for weak lines of growth, olive buff, profusely marked with black lines in irregular, exaggerated zigzag pattern. The aperture is bluish gray within, whitish toward the lip edge, which is very little thickened at the base, acute. Columellar area flat, nearly white, the columellar edge blunt, smooth; nearly straight. No operculum-stop within the baso-columellar angle.

Type Locality: Great River, Jamaica.

Remarks: Type, Academy of Natural Science of Philadelphia No. 153556; Vermeij--species possible a Neritina.

Nerita ampullacea

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 172.

Description: *Nerita testa perforata, ventricosa, supra depressa, apice obtuso; apertura larga.*

Type locality: Asiae paludosis.

Remarks: Now Helix.

Nerita anatolica

Recluz, M. C. A. 1842. *Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier.* 1842, p. 342.

Description: *Testa, subovata, violacea, tenue striata; spira brevissima, rotundato-obtusas; apice minimo, laterali; labio subobliquo, albido-caerulecente, infero dilatato, subconvexo, margine recto, acuto.*

Type locality: Smyrne, Alep., Sidon and Scio.

Remarks: Probably freshwater, Neritina.

Nerita angularis

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 187.

Description: *Testa opaca, conica, glabra virescens striis*

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. angularis, cont.

transversis, subtilissimis, spiralibus vero tribus in singulo anfractu elevatis, acutis. Anfractus quinque fere perpendiculares. Apertura rotundata, ad anfractum vicinum in angulum producta. Foramen vel umbilicus nullus. Faux calcarea. Striae spirales in quibusdam evanescunt.

Type locality: "In flumine Chinensi emporium Canton alluente."

Remarks: Vermeij--species questionably Nerita.

Nerita angulata

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

Nerita angulosa

Meuschen, F. C. 1787. Museum Geversianum, p. 274.

Remarks: Not acceptable.

Nerita angulosa

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 173.

Description: N. testa ventricoso-globosa, superne depresso-plana, subtus angulosa, olivaceo, maculis luteis parvulis obsoletis notata; anfractibus duobus inaequaliter rugatis; supremo dersos-concavo, apertura subrotundata, antice et postice angustata, superne et inferne dilatata; columella angusta, medio concava, superne callosa, antice tenuie denticulata, medio emarginata.

Type locality: Casam, province of Misamis, island of Mindanao, on stones in small stream.

Remarks: Probably not Nerita.

Nerita annulata

Reeve, L. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 18, species 78.

Description: Ner. testa suborbiculari, depressiuscula,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. annulata, cont.

crassa, spira leviter exserta, mucronata, anfractibus superne depresso-depressis, deinde obsolete angulatis, convexis, spiraliter annulato-constatis, apertura expansa, labro effuso, intus crenulato, columella superne obscure corrugata, subdentata; nigra, punctis paucis luteis variegata, inferne fasciis duabus albis latis cincta.

Type Locality: Unknown.

Nerita nodosaNerita anthracina

Busch, G. von dem. 1844. In R. A. Philippi. Abbildungen und beschreibungen neuer oder wenig gekannter conchylien, p. 85.

Description: N. testa globosa, opaca, rudi, striis transversis sc. et longitudinalibus irregularibus sculpta, nigra aut flavescente marmorata; labor intus denticulato; columella quadridentata, dentibus medianis minutis; labio concavo, granuloso.

Type Locality: Java.

Nerita antillarum

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3685.

Description: N. testa subglobosa nigra intus alba sulcata striataque: vertice obtuso, labio utroque rugoso denticulato.

Type Locality: "Habitat frequens ad insulas Antillas."

Nerita antiquata

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 106.

Description: N. testa subglobosa, crassiuscula, longitudinaliter rugosa, ex luteo fuscescente, nigro et albo obsolete maculata; spira brevissima, obtusa; apertura alba, fauce angustata, luteo-aurantia; labio convexo, rugoso et margine bidentato; labro rotundato, acuto, intus striato superne dentato.

Type locality: Philippine Islands.

--

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. antiquata, cont.Remarks: Philippine endemic, similar to N. polita.Nerita antiquata

Recluz, M. C. A. 1853. Description de quelques nerites et d'une natice. J. Conchyliol. 4, p. 317.

Description: N. testa subglobosa, crassiuscula, longitudinaliter rugosa, ex luteo-fuscescente, nigro et albo maculata, lineata, variegata vel fasciata; spira brevissima, obtusa, apertura alba, fauce angustata, luteo-aurantia; labio convexo, rugoso, 2-3 dentato; labro rotundato, intus stiato.

Type locality: Philippine Islands.

Nerita aperta

*Defrance, . 1825. Dict. Sci. Nat. 34, p. 479. [not seen]

Nerita apita

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 72.

Description: Ner. testa ventricosa-globosa, tenuissima, rugosa, subepidermide, olivacea fusco-violascente, lituris transversis albidis picta,; anfractibus quaternis superne depresso-planulatis; spira prominula, apice pulchre croceo, hyalino; apertura dilata; labio semilunaris, nigrescente externe zonato, atice albo, recto, integerrimo.

Type locality: Island of Negros, Philippine Islands.

Nerita arabica

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 5, species 20.

Description: Ner. testa compresse globosa, solida, spira vix exserta, anfractibus rotundatis, spiraliter linearis sulcatis, sulcis superficiariis, apertura incrassata et tenuiserrata, area columellari verrucoso-rugosa et dentata; olivaceo-viridi, roseo distincta, punctis et maculis nigris undique lessellata.

Type locality: Aden, Arabia. —

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita arachnoidea

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3674.

Description: *N. testa alba lineis rufescens reticulata apice nigricante: anfractibus convexis, umbilico subobtecto.* Chemn. Conch. 5, p. 188, f. 1915, 1916.

Type locality: Unknown.

Remarks: Check Cernohorsky.

Nerita argus

Recluz, M. C. A. 1841. *Description de quelques nouvelles especes de nerites vivantes.* Rev. Zool. Soc. Cuvier. 1841, p. 150.

Description: *Testa parva, semiglobosa, transversim tenue striata, subepidermide rufescens nigra, punctis croceis per series longitudinales obliquas pulchre picta, anfractu infimo superne suboblique depresso; apice retuso; apertura obovata, albido-viridescente; labio basi bidenticulato, supra rugoso et granoso; labro intus sulcato, superne bidentato, basi effundente.*

Type locality: Rio de Janiero.

Nerita arriaca

Bolten, J. F. 1798. *Museum Boltenianum*, Part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

Nerita articulata

Gould, A. A. 1847[1848]. [untitled]. Proc. Boston Soc. Nat. Hist. 2, p. 220.

Description: *Testa oblique semiglobosa, postice planulata, antice dilatata; spira prominula, anfr. 3, liris propinquis atro-purpureis, albo articulatis 25-30 cinctis, interspatiis latioribus lucidis carneolis desussatim striatis: apertura magna semilunari, labro acuto atropurpurco, allo albo creberrime sulcato firmato; fauce lutco; labio excavato, extrorsum plicato; columella scuta, sinuata, denticulis 3-4 armata.*

Type locality: Tavoy.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

N. articulata, cont.

Remarks: Replacement name for N. lineata Gmelin 1791, nec Muller 1774; see Mienis (1973), Argamon 4(1):21-22.

Nerita ascensionis

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3683.

Description: N. testa sulcis albis costisque albo naculatis sedecim, vertice prominulo, labio exteriore utrinque gla-
bro, interiore subconcavo flavicante dentato.

Type locality: Ascension Island

Nerita ascensionis deturpensis subsp. nov.

Vermeij, G. J. 1970. The Nerita ascensionis species complex in the south Atlantic (Gastropoda: Prosobranchia). Veliger 13(2), p. 135.

Description: Shell thin but strong, moderately depressed, greatest linear shell diameter (width) up to 30.4 mm; spire low, consisting of 3-4 whorls, often considerably corroded. External shell surface black, sculptured with 14-18 weak to moderately strong spiral ribs which are strongest near the suture; sometimes the ribs may disappear during ontogeny, so that the shell becomes smooth or nearly so in later stages of growth. Parietal area plane to somewhat concave, smooth to very weakly pustulose, bright light orange in color; edge of parietal area with 2 rather weak median teeth, the posterior one being the more strongly developed. Aperture proper (opening between outer lip and edge of parietal area) quite wide; outer lip somewhat thickened within, with about 16 very weak teeth, which are generally absent in young specimens; the posteriormost of these teeth is much the largest, but is still poorly developed in comparison to other members of the subgenus; no large anterior tooth on the outer lip. Operculum calcareous, thin; outer surface very slightly convex, brownish in color, with numerous darker-colored pustules.

Type locality: Praia da Atalaia, south coast of Ilha Fernando de Noronha.

Location of types: Holotype, Museum of Comparative Zoology No. 262991; paratypes, Museum of Comparative Zoology and Yale Peabody Museum.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita ascensionis trindadeensis subsp. nov.

Vermeij, G. J. 1970. The Nerita ascensionis species complex in the south Atlantic (Gastropoda: Prosobranchia). Veliger 13(2), p. 136.

Description: This subspecies is very similar to Nerita ascensionis deturpensis, but teeth on the outer lip are barely developed, and are in any case never as strong as in the populations from Fernando de Noronha. The parietal area is color a very pale light orange, being similar in this respect to typical N. a. ascensionis.

Type locality: Trindade Island.

Location of types: Holotype, Museum of Comparative Zoology No. 262990; paratypes, Museum of Comparative Zoology.

Nerita aspersa

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

Nerita asperulata

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 169.

Description: N. testa ovato-elliptica, striis intertextis granulosis asperulata, fusco-nigra, sub oculo et lumine, lineis nigris ramosis transversis intricatis notata; anfractibus 1 1/2, superne depresis; apice prominulo, obtuso; apertura rotundata, superne et inferne dilatata, luteo-fuscescente; labio plano-compresso, margine vix arcuato et tenue ruguloso.

Type locality: Psaguing, North Ilocos, Luzon, Philippine Islands (on stones in small river).

Nerita aterrima

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3679.

Description: N. testa crassa opaca globosa nigerrima lineis picta, intus alba: labio exteriore glabro, interiori tuberculato rugoso.

Type locality: Unknown.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. aterrima, cont.

Remarks: See Cernohorsky.

Nerita atra

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 188.

Description: *Nerita testa subulata, laevi, apertura antice posticeque sinuata.*

Type locality: From Spengleriano Museum.

Remarks: Now Strombus from Index Animalium 1.

Nerita atramentosa

Reeve, L. A. 1855. *Conchologia iconica: or, illustrations of the shells of molluscous animals* 9 (*Nerita*), plate 6, species 25.

Description: *Ner. testa semiglobosa, crassa, solida, spira vix exserta, anfractibus convexis, subangustis, laevibus vel tenu spiraliter stiatis, striis subundultais, labro intus incrassato, dente unico, columella plano-concava, margine minute denticulato; caeruleo-nigra, punctis albidis spiraliter lentiginosa.*

Type locality: Swan River, Australia.

Nerita atrata

Gmelin, J. F. 1791. *Systema naturae*, 13th edition, p. 3683.

Description: *N. testa aterrima glabra suprateniussime striata: labio utroque albo, exteriore intus subtiliter sulcato subdentato, interiore concavo rugoso tuberculato.*

Type locality: Atlantic Ocean [Recluz says Malaya].

Remarks: See Recluz, 1841, p. 74; see Cernohorsky.

Nerita atropurpurea

Recluz, M. C. A. 1841. *Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier.*
1841, p. 107.

Description: *Testa semiglobosa, supra depresso-plana, medio*

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. atropurpurea, cont.

angulosa, atropurpurea, transversim sulcata: spira brevisima, obtusa, ad latus oblique incurva; apertura subrotunda, intus angustata; labio posteriori rugoso, medio granoso, margine tridentato: dente prima validiore truncata: labro crenato, intus calloso, inermi, superne oblique dilatato spiram semi marginante.

Type locality: Apia ["iles Witi"].

Remarks: Vermeij--synonym of N. planospira.

Nerita aurantia

Recluz, M. C. A. 1842. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 73.

Description: Testa semiglobosa, acuta, sulcata: sulcis 35, superioribus latioribus inferioribus angustis, luteoaurantia, flammis cinereis obsoletis undulatisque picta; spira prominula, conico-acuta, dense et minutissime granulata; apertura lactea; labio convexiusculo rugoso ae tridentato; labro intus incrassato, sulcato, bidentato.

Type locality: Philippine Islands.

Remarks: "Tres-rare."

Nerita aurantiaca

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita auriculata

Wood, W. 1828. Index testaceologicus, p. 174.

Description: None given.

Type locality: New Holland.

Nerita aurita

Muller, O. F. 1774. Vermium terrestrium et fluviatilium, Vol. 2, p. 192.

Description: Nerita testa turrita, fusco-fasciata; anfractibus muricata; apertura ovata.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

N. aurita, cont.

Type locality: From ?museum.

Nerita aurora

Dunker, W. 1844. In R. A. Philippi. Abbildungen und beschreibungen neuer oder wenig bekannten conchylien, unter nithulf mehrere Deutscher Conchylionogen, herausgegeben von..., p. 87.

Description: N. testa crassa, ovato-semiglobosa, longitudinaliter plicata, transversim tenerime striata, nitida, alba, fasciis tribus croceis nigro variegatis picta; spira prominula acutiuscula; apertura semilunari aurantiaca; labio bidentato, obsolete rugoso, labro incrassato subacuto, intus tenerime crenulato. Operculum deest.

Type locality: Unknown.

Nerita australis

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3675.

Description: N. testa sub epidermide ex flavicante, virescente et caerulescente varia lineis angulatis viridibus picta: spirae anfractibus novem sulco dictinctis.

Type locality: Nova Seelandia.

Remarks: See Cernohorsky.

Nerita australis

Wood, W. 1828. Index testaceologicus, supplement, p. 25.

Description: None given.

Type locality: New Holland.

Nerita avellana

Recluz, M. C. A. 1842. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 76.

Description: Testa subglobosa, crassiuscula, roseo-violacea, maculis albis anterius nigro marginatis, sagittatis, cerebribus, 3 seu 4 faciata; sutura lactea, maculis nigris arcuatis alternis late cincta; spira

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. avellana, cont.

planulata; apertura alba, labio plano, angusto, margine arcuato et ruguloso.

Type locality: Manila [Manille].

Remarks: Described in the subgenus Clithon.

Nerita balteata

Reeve, L. A. 1855. *Conchologia iconica: or, illustrations of the shells of molluscous animals* 9 (*Nerita*), plate 6, species 28.

Description: Ner. testa ovato-subglobosa, solida, spira declivi, vix exserta, anfractibus subangustis, spiraliter liratis, liris tenuibus, funiculatis, labro intus incrassato, minute lirato, columella laevigata, superne corrugata, marbine obsolete denticulata; caerulescente-nigra, liris hic illic albo sparsim articulatis.

Type locality: Unknown.

Remarks: Species OK, see Cernohorsky.

Nerita beaniana

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 200.

Description: Ner. testa ovata, postice angustata, transversim crebre et tenuiter sulcata, nigra sive olivaceo-nigricante, fasciis croceis cincta; spira obtusa, decorticata; apertura alba, fauce luteo; labio compresso-plano, luteofuscescente, granulis nigris singato, margine laevissime arcuatim excavato, subedentulo; labro intus striis tenuisimis elongatis, nigris instructo.

Type locality: "Isle of Corregidor, bay of Manila."

Nerita beckii

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 275.

Description: Testa semi-globosa, subepidermide nigrescente rufo-fusca, immaculata; apice integro, planulato, angulo obsolete late cincto; columella nigro-caerulescente, margine recto, denticulato; labro rotundato, inus dilute

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. beckii, cont.

luteo-rufescente, superne libero, ad exteriorem arcuatim rotundato, acuto.

Type locality: "Mon cabinet et celui de madame veuve Dupont."

Remarks: Described in the subgenus Neritina.

Nerita bicanaliculata

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 171.

Description: N. testa ovata, postice truncato, dorso convexa, tenui, pellucida, fuscescente, lineis violaceo-nigris, crebre et eleganter reticulata; apice brevissimo, dorso interdum compresso plano; labio lato, subconvexo, margine arcuatim tenue denticulato; labro postice biauriculato; auriculis angustis, subtus canaliculatis.

Type locality: Island of Camiguin, found in small pond.

Remarks: "A fine Neritina, very distinct from all the auriculated species hitherto known."

Nerita bicolor

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 172.

Description: N. testa ventricoso-ovata, crassiuscula, sub sutura depressiusculo, strigis nigris et olivaceis longitudinalibus dense picta; apice eroso; apertura extus subquadrata exalbida; labio plano, superne calloso basi canaliculato, margine in medio arcuato, et obselete dentato; dente cardinali majori.

Type locality: "Agoo: in a mountain-stream."

Nerita bidens

Linnaeus, C. 1758. Systema naturae, p. 778.

Description: N. testa laevi, labio interiore bidentato.

Type locality: Unknown.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita bifasciata

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3685.

Description: *N. testa nigricante: faveis duabus canis,*
vertice albo.

Type locality: India.

Nerita bisecta

Reeve, L. A. 1855. *Conchologia iconica: or, illustrations*
of the shells of molluscous animals 9 (*Nerita*), plate 8,
species 29.

Description: *Ner. testa subglobosa, subexpansa, spira non*
exserta, anfractibus convexis, spiraliter costatis, costis
regularibus, subdistantibus, sulco undique bisectus, labro
intus effuso, minute crenulato, columella plana, vix den-
ticulata; caeruleo-nigra, area columellari vitellina, ni-
tente.

Type locality: West Africa.

Remarks: Vermeij--OK, Mauritius.

Nerita bizonalis

Lamarck, J. B. P. A. 1816. *Encyclopedie methodique his-*
toire naturelle des vers. Pl. 454, fig. 3a-b.

Description: None given.

Type locality: Not given.

Remarks: Lamarck (1838, v. 8, p. 606) listed this species
as synonym of *N. chamaelon*.

Nerita bourguignati

Recluz, M. C. A. 1852. *Recensement des nerites (sous-genre*
Neritine) de la France continentale. *J. Conchyliol.* 3,
p. 293.

Description: *N. testa parvula, transverse ovali, dorso con-*
vexa, per longum argute striata, fusco nigrita; anfracti-
bus tribus, superne in junioribus depresso-planiusculis,
. *in adultis convexiusculis; spira laterali, inclinata;*
apertura luteo-aurantia; labio plano, albido macula nigra
semper signata; labro tenui.--operculo croceo, apice

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

N. bourguignati, cont.
convexo.

Type locality: "Hab. riviere la Vaige, pres de la Bazouge
de Chemere, department de la Mayenne,..."

Nerita brevispina

Quoy, J. R. C., and J. P. Gaimard. 1834. Zoologie. In
Voyage de decouvertes de l'Astrolabe....p. 200.

Description: Nerita, testa semi-globosa, subepidermide
viridi-fuscescente zonata; ultimo anfractu superne angu-
lato, ad angulum spinis brevibus coronato; spira planius-
cula; apertura alba, labio denticulato.

Type locality: Havre Carteret, a la Nouvelle-Irlande.

Nerita bruguierei

Recluz, M. C. A. 1841. Description de quelques nouvelles
especes de nerites vivantes. Rev. Zool. Soc. Cuvier.
1841, p. 274.

Description: Testa ovato-subglobosa, tenui, rufo-fusca,
maculis nigris transversis seriatis fasciata; anfractu
infimo superne depresso, striato, nitidulo; apice convexo,
rugoso, integerrimo; columella valde plana, pallide
aurantia interdum albida, in medio obsolete crenulata et
vix arcuata; labro supra columellam arcuatim contorto.

Type locality: Les iles Philippines?

Nerita bullula

Reeve, L. A. 1855. Conchologia iconica: or, illustrations
of the shells of molluscous animals 9 (Nerita), plate 18,
species 76.

Description: Ner. testa ovata, spira obtusa, vix exserta,
anfractibus subangustis, laevibus vel spiraliter striatis,
solumella excavato-plana, subangusta, tenuissime obsolete
denticulata; carbonario-nigra.

Type locality: Unknown.

Nerita canalifera

*Buvignier, . 1852. [not seen] —

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita canalis

Wood, W. 1828. Index testaceologicus; or a catalogue of shells, . . . , Supplement, p. 26.

Description: None given.

Type locality: "South Sea Islands."

Nerita cancellata

*Hermann, J. 1781. Naturforscher, p. 56. [not seen]

Nerita candida

Wood, W. 1828. Index testaceologicus, p. 169.

Description: None given.

Type locality: Molluccas.

Nerita candidissima

*Salis, C. U. von. 1793. Reise Neapel, p. 380. [not seen]

Nerita canrena

Linnaeus, C. 1758. Systema naturae, p. 776.

Description: N. testa umbilicata laevi, spira submucronata, umbilido gibbo bifido.

Type locality: "O. Asiae."

Remarks: Vermeij--now Natica canrena from West Indies.

Nerita carbonaria

Philippi, R. A. 1844. Abbildungen und beschreibungen neuer oder wenig gekannter conchylien, . . . l, p. 84.

Description: N. testa ovata, laevi, nitida, lineis tenuis-simis impressis albo-punctatis sculpta; spira brevi; labro intus incrassato integerrimo; labio concavo, granulis duobus obsoletis notato; columella arcuata, medio denticulis minutis quinque ornata.

Type locality: Not given.

Remarks: Cernohorsky (1972) N. carbonaria syn. of N. morio; Rehder (1980) N. carbonaria nomen dubium until Philippi's

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

N. carbonaria, cont.
type located.

Nerita cardinalis

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 16.

Remarks: Nomen nudum, no valid standing.

Nerita cardinalis

LeGuillou, M. 1841. Description de quatorze Nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p.345.

Description: Testa ovata, purpureo-violacea; nitida, longitudinaliter subuplicata; anfractibus 4 1/2 superne angulatis; spira exsertiuscula, lineolis nigris picta, ad angulos brevissime spinosa; apertura dilatata, subquadrata; labio angusto, margine 4 denticulato.

Type locality: "Les iles Arrou" =Aru?

Nerita carinata

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable, check "Rejected Works."

Nerita cariosa

Wood, W. 1828. Index testaceologicus, supplement, p. 25.

Description: None given.

Type locality: Not given.

Remarks: Hawaiian freshwater nerite, Theodoxus cariosa; see Kay, p. 66.

Nerita carneae

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita celata

Recluz, M. C. A. 1845. Description de quelques nouvelles nerites fluviatiles, du cabinet de H. Cuming. Proc. Zool. Soc. London 13, p. 120.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. celata, cont.

Description: Ner. testa globoso-ovata, olivaceo-lutescente, postice pallidiore, striis intensioribus longitudinalibusque picta; anfractibus 1 1/2; infimo supra medium subanguloso; apertura albido-caerulescente pallido; columella piano-compressa, superne callosa, margine denticulata, medio breviter ac recte emarginata.

Type locality: Not given.

Nerita cerina

Colbeau, J.-A. J. 1859. Materiaux pour la fauna malacologique de Belgique 1, p. 11.

Description: Not given.

Type locality: Not given.

Remarks: Described as Nerita fluviatilis, L. variete Cerina; European Theodoxus.

Nerita cerostoma

*Troschel, F. H. 1852. Verzeichniss der durch Herrn Dr. v. Tschudi in Peru gesammelten Conchylien. Arch. Naturgesch. 18(1), p. 179. [not seen]

Type locality: Peru.

Remarks: Vermeij--no specimens seen, questionable identity; see von Martens for discussion.

Nerita ceylonensis

Recluz, M. C. A. 1851. Description de quelques coquilles nouvelles. J. Conchyliol. 2, p. 202.

Description: N. testa parvula orbiculato-subovata, ventricosa, nigerrima, longitudinaliter substriata; anfracitibus 1 1/2: ultmo subsutura spiratim depresso; apice minuto, punctiformi, convexo seu excavato; peritremate subcirculares; apertura semi-rotunda, albido-caerulescente; labio piano, antice subexcavato, acuto, edentulo, basi vix emarginato; labro acuto, superne dilatato, subformicato, ad suturam extus compresso.

Type locality: Ceylon (M. Morelet). —

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita chamaeleonLinnaeus, C. 1758. *Systema naturae*, p. 779.

Description: *N. testa sulcata sulcis viginti, labiis dentatis: interiore rugoso tuberculato.*

Type locality: ?Banda.

Remarks: Rumph. *Mus. t. 22, f. L. valvata 3 undulata.*

Nerita chemnitzii

Recluz, M. C. A. 1841. *Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier.*
1841, p. 103.

Description: *N. testa ovato-subgloboso, crassa, transversim striata, cinereo-lutescente, nigro trifasciata, flammis longitudinalibus albidis undato flexuosis, remotis fasciis secautibus ornata; spira brevissima, obtusa, lutescente; apertura ovata; labio lato, incrassato, convesiusculo, inferne vix rugoso, inaequaliter tridentato; labro intus striato, superne bidentato.*

Type locality: "Port-du-Roi-Georges, la Nouvelle-Holland et Timor."

Nerita chiloensis

*Molina, . 1810. *Sog. Star. Nat. Chili*, 2nd edition, p. 177. [not seen]

Remarks: May not be Nerita.

Nerita chlorina

Link, H. F. 1807. *Beschreibung der Naturalien-Sammlung der Universitat zu Rostock 3*, p. 141.

Remarks: No description.

Nerita chloroleuca

Philippi, A. R. 1848. *Testaceorum novorum centuria. Z. Malakozool.* 5, p. 14.

Description: *N. testa ovato-globosa, solida, alba, olivaceo maculata, transversim sulcata; sulcis 20-24; interstitiis rotundatis, costaeformibus; spira elevata; amfractibus rotundatis, ultimo superius depresso; apertura subrotunda,*

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. chloroleuca, cont.

patula, alba; labio concavo, extus et superius sulcato, centro granulato; margine columellari denticulis tribus minimis; labro intus denticulis circa 12, secundo superiore maximo.

Type locality: Not given.

Nerita chlorostoma

Lamarck, J. B. P. A. 1816. Encyclopedie methodique histoire naturelle des vers. Pl. 454, fig. 4a-b.

Description: None given.

Type locality: "Mon cabinet."

Remarks: Vermeij--see ascensionis.

Nerita chrysostoma

*Karsten, D. L. G. 1789. Museum Leskeanum 1, p. 290. [not seen]

Nerita chrysostoma

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

Nerita chrysostoma

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 104.

Description: N. testa ovato subglobosa, transversim sulcata, luteo-fulva nigro varia seu interdum subtifasciata, spira prominula, apice lutea, acuta; apertura ovata, flava; labio tridentato, superne incrassato, rugoso, labro subprolongato, intus tenue sulcata superne bidentata.

Type locality: Philippines.

Nerita cincta

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita cinerea

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita circumvoluta

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 173.

Description: N. testa globoso-subovata, dilute fulva in roseo vertente, punctis luteolis antice lineis nigris marginatis marmorata, zonis nigris remotis circumcincta; anfractibus duobus nigro-marginatis; supremo obtuso, dexorticato; apertura albida; labio plano, superne calloso, antice reguloso, et in medio arcuatim emarginato.

Type locality: Philippine Islands.

Remarks: Note "operculum of a Clithon" p. 173; IA2 added "Neritina (Recl), G. B. Sowerby, II, Thes. Conch. II(10) 1848, 527, Nerita."

Nerita columbaria

Recluz, M. C. A. 1845. Description de quelques nouvelles nerites fluviatile, du cabinet de H. Cuming, Esq. Proc. Zool. Soc. London 13, p. 121.

Description: Ner. testa ovato-globosa, laevissime striata; anfractu unico, refescente, viridescente pallido obsolete fasciata, maculis minimis luteis nigro seu roseo marginatis picta; columella callosa, subconvexo, denticulata, medio vix emarginata; labro antice subtruncato, basi et intus incrassato. Operculo crasso, extyerne viridulo, lutescente pallido undato, medio perlongum stria bipartito.

Type locality: Colombo, Ceylon.

Nerita comma-notata

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals (Nerita) plate 17, species 72.

Description: Ner. testa ovata, spira obtuse exserta, anfractibus subangustis, superne declivibus, spiraliter dense liratis et sulcatis, columella laevigata, leviter

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. comma-notata, cont.

denticulata; caeruleo-nigra, punctis albis commaeformibus aspersa.

Type locality: Unknown.

Nerita communis

Quoy, J. C. R., and J. P. Gaimard. 1834. Zoologie. In Voyage de decouvertes de l'Astrolabe...p. 195.

Description: Nerita, testa oblonga, laevigata, nigricante; lineis flavidis longitudinalibus, obliquis, angulatim flexuosis; apertura albicanti; labio luteo, denticulato.

Type locality: Vanikoro.

Nerita conoidea

*Lamarck, J. B. P. A. 1804. Ann. Mus. Nat. Hist., ser. 5, 26, p. 93. [not seen]

Nerita conspurcata

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 19.

Remarks: Nomen nudum, no valid standing.

Nerita constellata

Sowerby, G. B. 1905. Descriptions of new species of marine shells, chiefly from Ceylon. Ann. Nat. Hist. 16, p. 190.

Description: Testa oblique ovata, solidiuscula, laevis, purpurascens, maculis parvis albidis inaequalibus conspersa, spira brevissima, planulata fere immersa; anfractus ultimus rotunde convexus, concentrica obscurissime striatus, subobsolete spiratlier sulcatus; apertura semicircularis, intus laevis; labrum acutum; area columellari laevis, leviter convessa, ad marginem minute irregulariter denticulata

Type locality: Ceylon.

Nerita contorta

Muller, O. F. 1774. Vermium terrestrium et fluviatilium. Vol. 2, p. 187.

Description: Testa pyramidalis, squalida, opaca. Anfractus

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. contorta, cont.

vix quinque, convexi sibi invicem fere perpendiculariter impositi; quartus & quintus in ipso apice depresso, hicc lateribus tantum tres numerantur, & apex truncatus conspicator. Apertura circinnata est, ac centrum uit in antecedente perforatum.

Type locality: Not given.

Nerita cornea

*Grateloup, . 1828. [title unknown]. Bull. H. N. Soc. Linn. Bordeaux 2(10), p. 149. [not seen]

Remarks: Same as species below?

Nerita cornea

Linnaeus, C. 1758. Systema naturae, p. 777.

Description: N. testa obsolete striata, labiis edentulis.

Type locality: Not given.

Nerita corona

Linnaeus, C. 1758. Systema naturae, p. 777.

Description: N. testa anfractibus coronatis spinis, labiis edentulis.

Type locality: "Asiae fluviis" p. 777.

Remarks: Freshwater, Neritina?

Nerita corona

*deBlainville, H. D. 1824. [title unknown]. Dict. Sci. Nat. 32, p. 239. [not seen]

Nerita corrosula

Recluz, M. C. A. 1842. Description de plusieurs de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 177.

Description: Testa ovata, ventricosa, tenuiuscula, impolita, corrosiuscula, tenuiter striato-costata; spira laterali, prominula, rotundato-obtusa; apertura extus ovali et subcontinua; labio plano, albido-fuscescente, rugoso et bidentato; labro acuto, intus sulcato, superne subunidentato.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. corrosula, cont.

Type locality: New Guinea

Nerita costataGmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3684.Description: *N. testa intus flavicante subglobosa striis
crassioribus cincta: interstitiis niveis, labio utroque
dentato, exteriore exterius crenato, interiore subconvexo
rugoso tuberculato.*

Type locality: ..."insulas Indiae nicobaricas" (p. 3684).

Nerita crassaGould, A. A. 1852. *Mollusca and shells. In United States
Exploring Expedition during the years 1838.... 12*, p. 166.Description: *Testa solidissima, sub-globosa, longitrorsum
striata, spiraliter sulcato-costata (costis ad triginta
rotundatatis), fusco-cinerea, flammulis crebris angulatis
obliquis variegata: spira prominula, ad apicem lutea; an-
fractibus quator rotundatis; sutura impressa: apertura
parva, semicirculatis; labro crassissimo, anciso, crenu-
lato, intus sulcato, postice unidentato; labio angusto,
octo-plicato, plicis anticis bisectis; columella biden-
tata, dente antico minimo.*

Type locality: None given.

Remarks: Vermeij--possibly synonymous with N. undata from
Smithsonian notes.Nerita crepidulariaWood, W. 1828. *Index testaceologicus; supplement*, p. 25.

Description: None given.

Type locality: Not given.

Nerita croceaBolten, J. F. 1798. *Museum Boltenianum, Part 2*, p. 17.

Remarks: Nomen nudum, no valid standing.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita cruenta

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 19.

Remarks: Nomen nudum, no valid standing.

Nerita cruentata

Gmelin, J. F. Systema naturae, 13th edition, p. 3673.

Description: N. testa subglobosa alba sanguineo maculato et punctata apice obtuso intusque caerulescente: umbilico spirali.

Type locality: Not given.

Nerita cumingiana

Recluz, M. C. A. 1842. Description de neuf especes de nerites nouvelles, suivie d'observations sur les Nerita cornea et dubia. Rev. Zool. Soc. Cuvier. 1842, p. 74.

Description: Testa oblongo-acuto, crassiuscula, nigra, lineolis lutescentibus picta; anfractibus quinis, convexiusculis; spira prominenti, conico-acuta; apertura oblique obconica; labio vix convexo, superne calloso, aurantio, marigine ruguloso, in medio valde emerginato.

Type locality: Philippines and Moluccas.

Nerita cuprina

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 72.

Description: Ner. testa ventricoso-oblonga, subepidermide cupreo-micante nigra, lineolis angulato-flexuosis densissime picta; anfractibus 4-5? convexis, superne rotundatiusculis; apice deroso; labio plano, lutescente, in margine vix arcuato et denticulato, basi subemarginato; labro intus albido-caerulescente, ad marginem luteo-viridescente.

Type locality: Not given.

Nerita cuvieriana

Recluz, M. C. A. 1841. Description de quelques especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 338.

Description: Testa ventricoso-ovata, laevissime striata,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. cuvieriana, cont.

fulvo-olivacea, punctis maculisque dilute lutescentibus marmorata; anfractibus quaternis, superne depresso-depressis; spira exsertiuscula, apice saepius decorticata; apertura subrotunda, intus late semilunari; labio angusto, albo, posterius crocco, marine tenuem arcuato, crenulato, basi emerginato.

Type locality: Guadeloupe.

Nerita dacostae

Recluz, M. C. A. 1844. [untitled]. Proc. Zool. Soc. London 11, p. 199.

Description: Ner. testa subglobosa, supra medium vix angulosa, mutica, olivacea, maculis triangularibus albidis, antice acute nigris picta et fasciis pallidissime interdum obsoletis cincta; anfractibus quaternis plano-declivis; spira subconica, apice erosa; apertura subrotunda, intus alba et pallide caerulecente 3-4 fasciata; labio omnipresso, angusto, valde declivi, superne transversim calloso, margine crenato et in medio tenuiter arcuato; labro inferne dilatato, superne subfornicato.

Type locality: "Isle of Negros, in a mountain stream"
p. 199.

Remarks: Freshwater, Clithon.

Nerita debilis

*Dufo, H. 1840. Ann. Sci. Nat., Ser. 2, Zoology, 14, p. 365. [not seen]

Remarks: See Taylor (1968, p. 199) for specimen noted from Seychelles; Vermeij says Nerita, Meinis says not nerite.

Nerita deflorata

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable, check "Rejected Works."

Nerita deformis

*Sowerby, J. C. 1840. Trans. Geol. Soc. London 5,
. explanation to plate 57. [not seen] —

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita dendrites

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita deshayesii

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 104.

Description: N. testa globoso-acuta, crassa, nigerrima, interdum radiis croceis picta, transversim sulcata: costis rugosiusculis; spira exsertiuscula, subacuta; apertura coarctata; labio convexo, albo aut croceo rugoso et granuloso, margine dentibus tribus remotis validis; labro margine tenui nigro et luteo articulato, intus incrassato, striato et bidentato.

Type locality: "Les cotes de la Californie" (p. 104).

Nerita diadema

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 277.

Description: Testa subgloboso, lucida, laevissima, olacea, maculis triangularibus et oblongis, luteis, nigro marginatis picta; anfractibus 3-4; ultimo supra medium subanguloso, spinis brevibus, olivaceis, coronato; apertura caesa; labio angusto, crenulato.

Type locality: "Les iles de la mer de sud (Lesson)" (p. 277).

Remarks: Clithon.

Nerita diaphana

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita dilatata

Moquin-Tandon, A. 1855. Histoire naturelle des mollusques terrestres et fluviatiles de France, Vol. 2, p. 549.

Description: Not given.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. dilatata, cont.

Type locality: Not given.

Remarks: Form beta of Nerita fluviatilis, (Figs. 32, 33).Nerita dissimilisMuller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 184.Description: *Nerita testa subovato-acuminata, luteo-albescente; labro nigro.*

Type locality: [In Museo Spengleriano]

Nerita diversicolorMartyn, T. 1789. *Universal conchyologist*, Vol. 3, p. 108.

Remarks: Nonbinomial, rejected name.

Nerita doingiiRecluz, M. C. A. 1845. *Description de quelques nouvelles nerites fluviatile, du cabinet de H. Cuming, Esq. Proc. Zool. Soc. London 13, p. 121.*Description: *Ner. testa parva, globoso-ovata vel subgloboso, parum obliqua, tenuiuscula; anfractibus 1.5-2 oblique rugulosis, transversim et sub lente creberrime ac obsolete striolatis; apertura viridula; columella fere plana, margine denticulata et in medio subarcuata; labro tenui, semiciruclari.*

Type locality: Hanover Bay, North Australia.

Nerita doliolum*Schrank, F. v.P. 1796. *Sammel. Nat. Phys. Aufsat.*, p. 319.
[not seen]Nerita dombeyiRecluz, M. C. A. 1841. *Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 149.*Description: *Testa parvula, ovato-globosa, anterius dilatata, dilute olivacea, transversim striata et maculis nigris subtrifasciata, interdum unicolore; labio coarctato,*

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. dombeyi, cont.

plano posterius rugosiusculo, medio granuloso, margine in medio vix arcuato et subedentulo; labro dilatato intus albo, vix striato, ad basin labianculo calloso sotato.

Type locality: Bombay.

Remarks: Vermeij indicates that this species looks like N. insculpta, see Rehder and Smithsonian specimens.

Nerita donovani

Recluz, M. C. A. 1843. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 175.

Description: N. testa semiglobosa, tenue et dense striata, viridescente, lineis obliquis purpureo-nigrescentibus, equidistantibus undatis ornata; anfractibus tribus convexis, supra medium spinis incurvis armatis; apice deroso; apertura caeruleo-roseata, basi effuso, angulosa; labio angusto, superne calloso, basi concavo, margine subrecto denticulato; dente cardinali majori.

Type locality: Island of Guimaras, Philippines.

Nerita doreyana

Quoy, J. R. C., and J. P. Gaimard. 1834. Zoologie 3. Voyage des decouvertes des l'Astrolabe..., p. 190.

Description: Nerita, testa antice lata, transversim tenuissime striata, subtessellata, rubente, tribus fasciis albo et nigro variegatis cincta; apertura alba, levi, fusco marginata.

Type locality: New Guinea.

Nerita dubia

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3678.

Description: N. testa tenui pellucida ovata nitida obscurius flava nigro varia; labio exteriori acuto, interiori glaberrimo.

Type locality: Not given.

Remarks: Recluz (1842, p. 79) species 63 given as Nerita

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES^{*}N. dubia, cont.

dubia (Chemnitz), Gmelin provided Chemnitz 5, p. 193, as source, as did Recluz; Chemnitz not binomial.

Nerita dubia

Turton, W. H. 1932. Marine shells of Port Alfred, S. Africa. Oxford University Press, London, p. 202.

Description: The shape of the shell is helicoid; 1 minute nuclear whorl, the remaining 1 very large; the aperture is almost an exact semicircle. The surface is marked by fine spiral liens, and by what appears to be extremely fine lines of growth ar right angles to these. They are just visible between the sprial lines and do not cross them. The colour is greyish white, and the size 1.3 x 1.2 mm.

Type locality: Port Alfred, South Africa.

Remarks: Check Kilburn, Notes on South African shells for this species.

Nerita eburnea

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 20.

Remarks: Nomen nudum, no valid standing.

Nerita eburnea

*Hoeninghaus, F. W. 1831. J. Min. 1831, p. 139. [not seen]

Remarks: Index Animalium 2 listed as nomen nudum.

Nerita edentula

*Sassi, A. 1827. Giorn. Liguistico 1, ser. 5, p. 477. [not seen]

Nerita effusa

Muller, O. F. 1774. Vermium terrestrium et fluviatilium, Vol. 2, p. 175.

Description: Nerita testa umbilicata, vernrericosa, fasciata, supra deppressa, apice acuto; apertura effusa.

Type locality: "In Museo Moltkiano."

—
TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES*

Nerita electrina

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 17, species 70.

Description: Ner. testa suborbiculari, deppressa, spira subexserta, mucronata, anfractibus convexis, densa spiraliter liratis, interstitiis obsolete puncturatis, columella tenue denticualta; lutescente-aurantia, immaculata.

Type locality: Isle of Panay, Philippines.

Nerita ellyptica

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 346.

Description: Testa ventricosa-globosa; basi obstuse acuta, longitudinaliter sublente striata; fusco nigricante, maculis triangularibus flavescentibus picta; anfractibus tribus, superne spinis nigris rectis immaculatis armatis; spira retuso-erosa; apertura rotundata; dilatata; labio angusto, plano, margine arcuato, superne inaequaliter crenato.

Type locality: Nuku-hiva.

Nerita elongata

*Philippi, R. A. 1844. Enum. Moll. Siciliae, vol. 2, p. 139. [not seen]

Nerita erubescens

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 12, species 54.

Description: Ner. testa globoso-turbinata, crassa, spira prominula, anfractibus convexis, spiraliter liratis et striatis, labro dente acuto laterali, columella leviter corrugata et dentata; pallide rufa, aut carneo-alba, punctis rufescentibus aspersa et fasciata.

Type locality: Not given.

• Remarks: May be synonmyzed with N. reticulata.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES*Nerita essingtoni

Recluz, M. C. A. 1842. Description de neuf especes de nerites nouvelles, suivie d'observations sur les Nerita cornea et dubia. Rev. Zool. Soc. Cuvier. 1842, p. 73.

Description: Testa semiglobosa, acuta, dense striata, roseo-violaceous, nigro maculata et subflammulata; spira prominula, conico-acuta; apertura alba, fauce flavidante; labio plano, ruguloso, bidentato; labro intus sulcato et unidentato.

Type locality: "La Nouvelle-Hollande, sur le detroit de Torres, a la pointe Essington" (p. 74).

Remarks: See Cernohorsky 3.

Nerita exarata

*Pfeiffer, L. 1840. Arch. Naturg. 6, p. 255. [not seen]

Nerita excavata

Sowerby, G. B. 1883. Thesaurus conchyliorum, or monographs of genera of shells, Vol. 5, p. 109.

Description: Oblique ovato-globosa, nigra albo variegata, costis validis, subcomplanatis approximatis lirata, spira subproducta obtusa; apertura flavidia, nigro marginata, profunda; area columellari granulata, declivi, excavata, dentibus marginalibus tumidis, acutis; labio externo intus sublirato, supra infraque acute bidentato; margine externo nigro, crenulato.

Type locality: Not given.

Remarks: Description from Sowerby, 1887, p. 109, species 25.

Nerita exuvia

Linnaeus, C. 1758. Systema naturae, p. 779.

Description: N. testa sulcata, labiis dentatis; interiore tuberculato.

Type locality: "O. Americae & Asiae."

Nerita fascia

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 19.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES^aN. fascia, cont.

Remarks: Nomen nudum, no valid standing.

Nerita fasciata

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 182.

Description: *Nerita testa ovato-ventricosa, alba, fasciis tribus rubris, splendidis.*

Type locality: "In Italia & Saxonia."

Remarks: Four other N. fasciata species described: Meuschen, 1787, p. 268, 272, 274, all probably invalid, see "Rejected Works;" also Martyn, 1789, fig. 110, nonbimomial, see "Rejected Works;" also Revier, 1804, see IA2.

Nerita fibrata

Colbeau, J.-A.-J. 1859. *Materiaux pour la fauna malacologique de Belgique* 1, p. 11.

Description: Not given.

Type locality: Not given.

Remarks: Described as Nerita fluviatilis, L. variete Fibrata; European Theodoxus.

Nerita filosa

Reeve, L. A. 1855. *Conchologia iconica; or illustrations of the shells of molluscous animals* 9 (*Nerita*), plate 10, species 48.

Description: *Ner. testa globoso-turbinata, spira exserta, mucronata, anfractibus superne declivibus, deinde convexis, liris minute bisectes spiraliter filosis, labro intus incrassato, minute crenulato, dente laterali acuto utrinque armato, columella laevigata, denticulata; ferrugineo-spadicea, liris eximie caeruleo-nigricantibus.*

Type locality: Not given.

Nerita flammea

Gmelin, J. F. 1791. *Systema naturae*, 13th edition, p. 3685.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. flammea, cont.

Description: *N. testa subglososa confertim transverse striata alba; radiis undalatis purpurascentibus, labio utroque dentato.*

Type locality: "Insulas Americae mediae" (p. 3685).

Nerita flammulata

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 180.

Description: *Testa ovato-semiglobosa, ventricosa, tenue sulcata, striataque, albido-cinerascente, flammulis undulatis nigris picta; spira brevi, convexo-depressa, obtusata, lutea; labio plano, rugoso, lutescente, in margine bidenticulato; labro crenulato, intus striato superne obsolete unidentato.*

Type locality: "Les iles de la mer du Sud, les Moluques?"

Remarks: Vermeij--good name for species from mangroves from Palau or Singapore.

Nerita flava

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

Nerita flavescens

Dillwyn, L. W. 1817. A descriptive catalogue of recent shells, arranged according to the Linnaean method..., Vol. 2, p. 992.

Description: Shell smooth, yellowish variegated with white, and three black bands; inner lip two-toothed, and the outer lip slightly striated.

Type locality: Nicobar Islands.

Nerita fluctifera

Meuschen, F. C. 1787. Museum Geversianum, p. 266.

Remarks: Not acceptable.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita fluviatilis

Linnaeus, C. 1758. *Systema naturae*, p. 777.

Description: *N. testa rugosa, labiis edentulis.*

Type locality: "Habitat in Europae cataractis."

Remarks: Type of Theodoxus, European freshwater form.

Nerita fluviatilis var. trifasciata

Colbeau, J.-A.-J. 1859. *Materiaux pour la faune malacologique de Belgique*, pl. 2.

Description: Plusieurs des fils ont abondonne leur place normale et se sont resserres en certains endroits de maniere a former bande.

Nerita fluviatilis var. undulata

Colbeau, J.-A.-J. 1859. *Materiaux pour la faune malacologique de Belgique*, plate 2.

Description: Le reseau compose seulement de fils longitudinaux, les autres manquant.

Type locality: Not given.

Nerita fluviatilis var. vermiculata

Colbeau, J.-A.-J. 1859. *Materiaux pour la faune malacologique de Belgique*, plate 2.

Description: Le reseau dechire obliquement dans certaines parties. On a quelque peine a compendre dans cette variete, la formation des petites lineoles vermiculees.

Type locality: Not given.

Nerita fontinalis

*Braud, C. P. 1815. *Hist. Coq. Paris*, p. 196. [not seen]

Nerita forskaelii

Recluz, M. C. A. 1841. *Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier.*
1841, p. 147.

Description: *Testa semi-globosa, rudi, transversim obsolete*

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. forskaelii, cont.

sulcata, cornea; spira exsertiuscula, subacuta, labio
plano-concavo, laevissimo, margine dentibus quaternis
obsoletis; labro intus striato.

Type locality: Red Sea.

Remarks: Vermeij says now N. sanguinolenta, Red Sea
equivalent of N. albicilla.

Nerita fragum

Reeve, L. A. 1855. Conchologia iconica: or, illustrations
of the shells of molluscous animals 9 (Nerita), plate 9,
species 41.

Description: Ner. testa globosa, spira vix exserta, anfrac-
tibus rotundatis, spiraliter tenue confertiim costata,
labro intus incrassato, tenue lirato, columella declivi-
excavata, granata et corrugata; caeruleo-nigra, strigis
perpaucis luteis erosionis notata, area columellari sangu-
neo-tincta.

Type locality: Not given.

Remarks: Characteristic blood spot on the granulated area
of the columella may make this species synonymous with N.
reticulata.

Nerita fuliginata

Reeve, L. A. 1855. Conchologia iconica: or, illustrations
of the shells of molluscous animals 9 (Nerita, plate 11,
species 50).

Description: Ner. testa ovato-semiglobosa, spira subex-
serta, anfractibus convexis, spiraliter dense linearis-
sulcatis, columella plano-excavata, tenue denticulata;
carnea lineis rufis undulatis confertis variegata, in-
tense nigro undique plus minus inquinata.

Type locality: Not given.

Nerita fulgurans

Gmelin, J. F. 1791. Systema naturae, 13th edition,
p. 3685.

Description: N. testa subglobosa confertim transverse
striata nigerrima; radiis ochroleucis; labio utroque

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. fulgurans, cont.

subdenticulato; interiore medio tuberculato.

Type locality: "Insulas Americae."

Remarks: Valid West Indian species.

Nerita fulminea

Gmelin, J. R. 1791. Systema naturae, 13th edition,
p. 3672.

Description: N. testa subglobosa lineis angulatis spaciceis
picta, lobo explanato.

Type locality: Africa.

Nerita funiculata

Menke, K. T. 1850. Conchylien von Mazatlan, mit kritischen
Anmerkungen. Z. Malakozool. 7(11), p. 169.

Description: N. testa-ovato hemisphaerica, fusco-cinerea,
nigro varia vel subzonata; spira convexa; anfractu ultimo
liris confertis, lamellis brevibus confertissimis imbrica-
tis asperis concto; labio granoso; margine columellari
medio 2- ad 3-dentato; labro intus dentato-lirato.

Type locality: Not given.

Remarks: Eastern Pacific valid species.

Nerita funiculata

Reeve, L. A. 1855. Conchologia iconica: or, illustrations
of the shells of molluscous animals 9(Nerita), plate 2,
species 9.

Description: Ner. testa ovata, solida, spira subexserta,
anfracitbus rotundatis, liris tenuibus confertis spirali-
ter funiculatis, apertura subincrassata, vix serrata, sub-
ampliter expansa, area columellari promiscue orrugata,
dentata; purpurascente-nigra, inter liras pallida.

Type locality: Not given.

Remarks: Eastern Pacific valid species?

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES*Nerita fuscata

Menke, K. T. 1829. Berzeichniss der anfelnlichen Conchylie-Sammlung der Freiherrn von der Malsburg, p. 15.

Description: Testa transversim dense sulcata, sulcis planiusculis, cinereo-nigra; spira brevi; labio dentato, supra verrucoso, macula rufa notato.

Type locality: Not given.

Remarks: Species 319 in list, one of two Nerita with descriptive information; Vermeij, possible synonym of N. funiculata.

Nerita fuscofasciata

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita gaimardi

Souleyet, M. 1842. Description de cinq nouvelles especes de nierites fluviales provenant du voyage de la Bonite. Rev. Zool. Soc. Cuvier. 1842, p. 269.

Description: Nerita, testa ovata-obtusa, viridescente, lineolis nigris longitudinalibus remotis undulatisque adspersa; spira prominula, rotunato-obtusa; labio albido lutescente, margine denticulato.

Type locality: Tourane, Cochinchine.

Nerita gemmata

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

Nerita gemmulata

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of shells of molluscous animals 9 (Nerita), plate 9, species 40.

Description: Ner. testa ovato-semiglobosa, tenuicula, spira subexserta, apice mucronata, anfractibus convexis, spiraliter linearis-liratis, liris crenulato-gemmatis, columella plano-excavata, margine integro; griseo-albida, liris caeruleo-nigro et opaco-albo tenue articulatis, apice

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

N. gemmulata, cont.
lutescente.

Type locality: Not given.

Nerita genuana

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of shells of molluscous animals 9 (Nerita), plate 18, species 77.

Description: Ner. testa orbiculari-ovata, depressiuscula, spira impresso-plana, anfractibus subangustis, versus aperturam latioribus, spiraliter tenuiliratis, labro intus fortiter crenao, columella declivi-excavata, profuse granulata, edentulata; intese nigra, zonis suabus latis luteis aut roseis cingulata.

Type locality: Not given.

Nerita georgina

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 180.

Description: Testa ovato elliptica, convexa, transversim tenue sulcata, albido-cinerea, maculis nigris obsoletis trifasciata; spira brevissima, obtusa, laterali; apertura ovata, intus coaretata; labio plano laevi, albido lutescenti, margine olsolete tridentato; labro integro, superne depresso, annulo luteo picta, subtus albo calloso integerrimo.

Type locality: "La Nouvelle-Hollande du port du roi Georges, d'ou ele a été rapportee par MM. Q. et Gaimard, selon le Museum de Paris" p. 180.

Remarks: Vermeij--questionable valid species, see von Martens discussion.

Nerita glabra

Wood, W. 1828. Index testaceologicus, Supplement, p. 26.

Description: None given.

Type locality: Not given.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

Nerita glaberrissima

*Brown, T. 1818. Mem. Werner. Nat. Hist. Soc. 2, p. 532.
[not seen]

Nerita glaucina

Linnaeus, C. 1758. Systema naturae, p. 776.

Description: N. testa umbilicata laevi, spira obtusiuscula, umbilico semiclauso; lobo gibbo dicolore.

Type locality: "O. Europaeo."

Nerita granulata

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 18, species 79.

Description: Ner. testa depresso-globosa, solidiuscula, spira plano-impressa, anfractibus spiraliter liratis, apertura expansa, labro intus crenato, columella profuse granulata; caeruleo-nigra, columella pallidissime aurantio-tincta.

Type locality: Not given.

Nerita granulosa

Meuschen, F. C. 1787. Museum Geversianum, p. 274.

Remarks: Not acceptable.

Nerita grayana

Recluz, M. C. A. 1844. [untitled]. Proc. Zool. Soc. London 11, p. 200.

Description: Ner. testa ovato-globosa, dorso-oblique conoidea, transversim sulcata; sulcis costis latioribus, striis longitudinalibus creberrimis sculpta; anfractibus sordide nigro-violaceis, obsolete albo maculatis; spira prominula, conico-depressa, acuta; apertura pallide fuscescente; labio planulato, margine tridentato, supra rugis confertis valde impresso; labro inut regulariter sulcato, superne unidentato.

Type locality: "Port Curimao, province of North Ilocos, isle of Luzon; on the rocks at low water." H. Cuming.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. grayana, cont.

Remarks: Valid species.

Nerita grisea

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita grisea

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 5, species 23.

Description: Ner. testa oblonga-ovata, spira subexserta, anfractibus superne concavo-impressis, deinde rotundatis, dense spiraliter sulcatis, apertura parum imcrassata, subserrata, area columellari subobsolete corrugato-lirata, denticulata; albida, griseo maculata et obscure trifasciata, apice pallide lutescente.

Type locality: Island of Samar, Philippines; Cuming.

Remarks: Vermeij--distinct species.

Nerita grossa

Linnaeus, C. 1758. Systema naturae, p. 779.

Description: N. testa sulcata, labiis dentatis; interiore convexo rugoso.

Type locality: O. Asiae.

Remarks: Valid, Indonesian endemic.

Nerita guamensis

Quoy, J. R. C., and J. P. Gaimard. 1834. Zoologie 3. Voyage des decouvertes des l'Astrolabe..., p. 191.

Description: Nerita, testa transversim striata, alba bel lutea, fasciis quaternis albo-rubris cincta; spira prominenti; apertura ampla, levi et alba.

Type locality: Guam.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES.Nerita querini

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 314.

Description: Testa ovato-semi-globosa solidiuscula, transversim striata, roseo-violacea, zonis nigrescentibus 3-4 fasciata, maculis minimis albis undique aspersa; spira laterali, brevi, obtusissima; apertura rotundata, albida, intus semi-lunari; labio plano, in medio vix arcuato, obsolete crenulato.

Type locality: Sumatra.

Nerita guttata

Meuschen, F. C. 1787. Museum Geversianum, p. 266.

Remarks: Not acceptable.

Nerita guttata

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 316.

Description: Testa ovata, nigerrima, guttulis lacteis vix perspicuis notato; anfractibus tribus convexiusculis; suturis superficialibus; spira exsertiuscula, rotundato-convexa; spertura cinereo-lutescente, labio angusto, piano, margine rectiusculo, tenuiter crenato.

Type locality: New Guinea.

Nerita haneti

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 181.

Description: Testa semi-globosa, tenui, nigerrima, nitidula, transversim obsolete striata; striis raris, remotis; spira brevi, convexiuscula; apertura rotundata, sordide aurantia; labio angusto, compresso, declivi, inmedio margine obsolete bidentato; labro intus et extus integerrimo.

Type locality: Marquesas Islands.

Remarks: See Rehder (1980, p. 22) note; on Easter Island, a synonymy of morio or argus says Vermeij; Rehder, p. 23.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

Nerita haustrum

Reeve, L. A. 1855. *Conchologia iconica: or, illustrations of the shells of molluscous animals* 9 (Nerita), plate 7, species 34.

Description: Ner. testa ovato-glogosa, spira vix exserta, anfractibus leviter linearis-sulcatis, labro intus lirato, columella concavo-planata, margine obsolete denticulata; olivaceo-nigra, lutescente-albo hic illuc interrupta, columella pallidissime roseo-tincta.

Type locality: Red Sea.

Nerita helicinoides

Reeve, L. A. 1855. *Conchologia iconica: or, illustrations of the shells of molluscous animals* 9 (Nerita), plate 18, species 80.

Description: Ner. testa ovato-turbanata, spira subampla, exserta, anfractibus juxta spiram depressiusculis, rotundatis, subangustis, versis aperturam contractis, sprialiter tenuicostatis, apertura parva, labro effuso, columella declivi, obscure denticualta; lutescente, maculis nigris et roseis alternatim fasciata.

Type locality: Not given.

Nerita helicinoides var. laevilabris

Pilsbry, H. A. 1895. *Catalogue of the marine mollusks of Japan*. Detroit. p. 177.

Description: Shell obliquely globose-turbinate, multilirate, lirae 18-20 on body-whorl, the lower 3 or 4 nearly obsolete; buff or whitish, encircled by three bands of black variegation alternating with similar rose-dotted zones; on some specimens the black replaced by rose. Spire elevated; last whorl descending, appressed at suture, concave below it. Aperture oblique, outer lip prolonged above, bevelled to an edge inside, margin yellow with dark dots, within this strongly thickened by a white smooth callus which is generally subtruncate or most weakly unidentate above. Area flat, smooth, yellow, the columellar edge with two slight submedian notches.

Type locality: Yaeyama, Okinawa, Japan.

Remarks: Synonym of N. guamensis (Vermeij). —

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita helicinoides var. tristis

Pilsbry, H. A. 1901. New Japanese marine, land and fresh-water Mollusca. Proc. Acad. Nat. Sci. Philadelphia, p. 397.

Description: Shell black with some white spots along the basal margin, and sometimes a few angular pink and white spots elsewhere. Columella three-notched in the middle; area smooth, yellow-tinted in the middle; lip-rib weakly crenulate, a small denticle near its upper end.

Type locality: Loo Choo Islands.

Remarks: Types Acad. Nat. Sci. Phil. Type No. 80,406.

Nerita hepatica

*Gray, J. E. 1821. London Med. Reposit. 15, p. 239. [not seen]

Remarks: Taken from Index Animalium 2; indicated as Nerita Syncera, currently a genus in Assmineidae.

Nerita hians

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita hieroglyphica

Dillwyn, L. W. 1817. A descriptive catalogue of recent shells arranged according th the Linnaean method, p. 995.

Description: Shell sub-globular, white, with blackish undulated streaks; inner lip toothed and crenulated.

Type locality: East China Seas.

Remarks: Species from Cheminitz 5, p. 322.

Nerita hilleana

Dunker, G. 1871. Mollusca nova Musei Godeffroy Hamburgensis. Malakozool. Blatt. 18, p. 167.

Description: Testa ovato-semiglobosa, transversim regulariter denseque contata, striis incrementi imbricatis et undulatis obducta, spira valde obtusa, perparum prominens; columella media sinuata plerumque indistincte

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. hilleiana, cont.

quadridentata, spetum paullo concavum, superne callosum, laevissimum; labrum crenatum, interne incrassatum et laeve; costae nigerrimae passim albo tesselatae, interstitia vel solei cinereo-virescentes et pellucentes, color columellae, speti et labri pallide sulphureus. Latit. vel diam. transv. speciminum majorum 16 mill., alt. 11 mill.

Type locality: Samoa.

Nerita hindsii

Recluz, M. C. A. 1844. [untitled]. Proc. Zool. Soc. London 11, p. 201.

Description: Ner. testa semiglobosa, solida, cinerea sive albida, nigro obscure articulata, transversim laeviter sulcata, striis longitudinalibus undulatis sulcis interdum decussantibus insculpta; spira parva, convexo-depressa, vix exserta; apertura dilute straminea; labio angusto, compresso-subconcavo, rugoso et granuloso, margine in medio 2-4 denticulato; labro intus valde calloso et longe sulcato. Operculum pallide cine-eo-fuscescens, granulosum, obsolete biangulosum, albido-viridescens, antice angulum planissimum circumdate. Dente apicali plano, superne laeviter incrassato, brevi, truncato, basi macula fusca motato; infimo arcuato, austriato, postice parum dilatato.

Type locality: "Ilo-Ilo, isle of Panay; under stones at low water." H. Cuming.

Nerita histrio

Linnaeus, C. 1758. Systema naturae, p. 778.

Description: N. testa sulcata transverscque striata, labio interiore dentato.

Type locality: Not given.

Remarks: Indonesia and Papua New Guinea endemic--Vermeij.

Nerita imperfecta

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES*Nerita incerta

Busch, G. von dem. 1844. In R. A. Philippi. Abbildungen und beschreibungen neuer oder wenig gekannter conchylien... . p. 85.

Description: N. testa ovato-globosa, nigra; anteriore parte opaca, posteriore nitida, punctis lacteis adspersa; labio intus minutim denticulato; columella medio denticulis quinque ornata; labio concavo, laevi.

Type locality: Java.

Remarks: See Meinis (1970); valid species from central Indo-Pacific (PNG, Solomon Islands)--Vermeij.

Nerita incurva

*Martens, E. von. 1887. Die Gattungen Nerita und Neritopsis, p. 50. [not seen]

Type locality: Querimba-Inseln.

Remarks: Information from illustration legend: Tafel 8, fig. 14-16.

Nerita indica

Souleyet, M. 1842. Description de cinq nouvelles especes de nerites fluviales provenant du voyage de la Bonite. Rev. Zool. Soc. Cuvier. 1842, p. 269.

Description: Testa ovata, transversa, striata, punctis seu lituris longitudinaliter seriatis picta; spira minima, oblique incurva, adpressa; apertura ovato-orbiculari, continua, intus rufo-unizonata; labio planiusculo, nigro, antice arcuato et denticulato.

Type locality: "Le Gange" [=Ganges?].

Nerita insculpta

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 152.

Description: Testa ovato-elliptica, transversim late saccatis, longitudinaliter striis creberrimis pulchre insculpta, sordide lutea; costis angustis lineis transversis nigris et albis maculata; spira brevissima, obtusa, laterali; apertura ovata, fauce coarctata; labio lato,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. insculpta, cont.

convexusculo, laevi, inferne callo notato, margine tenue arcuato, subdentato seu edentulo; labro crenulato intus calloso integro.

Type locality Timor.

Remarks: Valid species.

Nerita interrupta

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 173.

Description: N. testa ovato-semigloboso, apice valde corrossa et truncata, olivacea, lineolis luteis interdum interruptis creberrimis et fasciis latiusculis luteis transversis picta; apertura extus semi-ovali, intus luteo; labio subsemilunari, plano-concavo, margine in medio arcuato, acuto, dente cardinali exerto.

Type locality: "Yba, province of Zambales, ilse of Lucon: on stones in a rapid river."

Nerita intexta

Recluz, M. C. A. 1841. Description de quelques especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 340.

Description: Testa ovato-elliptica, tenui striata, lineolis nigris tenuissimis, crebris intertexta, interstitiis candidis, anfractibus tribus, superne depresso; spira exsertiuscula, papillata; apertura lutea, labio piano, acuto, albo.

Type locality: Brescia and Verone, in fresh water.

Remarks: Neritina.

Nerita intricata

Wood, W. 1828. Index testaceologicus, Supplement, p. 25.

Description: None given.

Type locality: Britian, etc.

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESNerita islandica

Gmelin, J. F. 1791. *Systema naturae*, 13th edition, p. 3675.

Description: *N. testa globoso-subacuta crassiore; anfractibus quatuor per sulcos profundissimos discretis.*

Type locality: "Habitat in Oceano septentrionali, an hujus tribus?"

Nerita jaculator

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 185.

Description: *Nerita testa oblonga, cornea, vertice acuto.*

Type locality: "In lacubus & rivis, vulgaris."

Nerita japonica

Dunker, G. 1860. *Neue japonische Mollusken. Malakozool. Blatter 6*, p. 233.

Description: *N. testa ovata, tenuicula, luteo et atro variegata; spira subelevata; apice et in pullis erosa; labro paullulum incrassato et columella laevigatis; labro subconcavo; operculo granoso.*

Type locality: Japan.

Remarks: Valid Japanese endemic species.

Nerita jovis

Recluz, M. C. A. 1843. [untitled]. *Proc. Zool. Soc. London 11*, p. 72.

Description: *Ner. testa ventricoso-ovata, nigra, lineolis angulato-flexuosis fulgurantibusve albis picta, punctataque; anfractibus quinis convexis; spira conico-depressa, nigro-violascente, albo punctata, acuta; apertura albo-virescente; labio plano, margine in medio vix arcuato adenticulato.*

Type locality: Not given.

Nerita junghuhni

Schepman, M. M. 1901. Note VIII. Description of a new

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

N. junghuhni, cont.

species of Nerita. Notes Leyden Mus. 13, p. 63.

Description: Shell semiglobose, solid, nearly smooth, with slight radiating striae; colour dark yellowish-grey, with indistinct white spots, forming spiral rows, some of which alternate with black spots, giving to the shell a mottled appearance; whorls about 2 1/2, spire scarcely exserted, more or less eroded, showing a yellow layer; a few younger specimens are covered by a yellowish epidermis; suture distinctly descending in front. Aperture small, greyish blue; lip with a thick white callosity with a blunt tooth, scarcely visible in many specimens. Columella with a broad but short tooth in the upper part, slightly excavated in the middle, where it bears 2 or 3 denticles, then slightly projecting again; columellar area white, smooth. Operculum flesh-coloured, with a radiating groove, dividing the external surface in two parts, of which the marginal part shows many small granules, which are less crowded and larger on the central portion; internal surface smooth, darker coloured, greyish near the nucleus; rib flattened, nearly white.

Type locality: Java.

Remarks: See Meinis (1970) and olivaria.

Nerita keraudrenii

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 346.

Description: Testa ventricosa-globosa; basi obtuse acuta, longitudinaliter sublente striata; fusco nigricante, maculis triangularibus flavescentibus picta; anfractibus tribus, superne spinis nigris rectis immaculatis armatis; spira retuso-erosa; apertura rotundata; dilatata; labio angusto, plano, margine arucato, superne inaequaliter crenato.

Type locality: Noukahiva.

Remarks: Remarkable operculum with small granulations.

Nerita knorri

Recluz, M. C. A. 1841. Description de quelques nouvelles especes der nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 274.

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. knorri, cont.

Description: T. ovata, crassiuscula, tenuiter striata, nigrante, immaculata; spira nulla; apice rotundato, integro; labio lato, anterius plano, posterius convexo, caerulecente, subarcuato, crenulato; labro superne arcuato, intus intense croceo, linea angusta, pallida, maringato.

Type locality: Madagascar.

Nerita labeo

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, vol. 2, p. 180.

Description: Nerita testa oblonga, umbilicata, fusca, punctis convexis striata, labro albo, dilatato.

Type locality: "In Museo Moltkiano."

Nerita lactaria

*Linnaeus, C. 1771. *Mantissa.....*, vol. 2, p. 551. [not seen]

Remarks: Synonym of N. plicata, see Adams and Leloup.

Nerita lacustris

Linnaeus, C. 1758. *Systema naturae*, p. 778.

Description: N. testa laeviuscula cornea, apice exquisito, labiis edentulis.

Type locality: "In Europae lacubus."

Remarks: Probably not a Nerita.

Nerita laevida

*Laskey, J. 1811. *Mem. Werner Nat. Hist. Soc.* 1, p. 409. [not seen]

Nerita laevilirata

Sowerby, G. B. 1914. Descriptions of fifteen new Japanese marine Mollusca. *Ann. Mag. Nat. Hist.*, Ser. 8, 14, p. 36.

Description: Testa suboblique globosa, pallidula, ferrugineo dense marmorata, maculata, et strigata; spira parva,

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIES

N. laevilirata, cont.

depresso conica; anfractus 3, planulati, spiraliter lirati; anfractus ultimus inflatus, liris rotundatis parum elevatis latis et angustis alternatim instructus, oblique irregulariter striatus; apertura parviuscula, alba, intus breviter plicato-dentata; labrum ampliter effusum, leviter concabum, ad marginem acutum; columella late callosa, medio panei pustulato, margo dextro pauci plicata, sinistro irregulariter rugose plicata.

Type locality: Oshima, Japan.

Nerita largillierti

Philippi, A. R. 1848. Testaceorum novarum centuria. Z. Malakozool. 5, p. 160.

Description: N. testa ovato-globosa, sulcis transversis superficialibus exarata, albida, maculis magnis nigricantibus picta; sulcis in anfr. ultimo 20-24; anfractibus convexis, ultimo denique superius depresso; spira elevata, obtusa; columella tridentata, dente superiore maximo, duabus aliis minutis; labro intus denticulis circa 15 munito, duabus superioribus majoribus, labio concavo, granulis notato, centro aurantio.

Type locality: Gabon in Guinea.

Nerita larva

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3679.

Description: N. testa laevi subglobosa alba faveis ex fusco luteis varia; vertice obtuso, labio subdenticulato.

Type locality: "Habitat rarissima ad Amboinam, testa mediae magnitudinis."

Remarks: See Cernohorsky.

Nerita latilinea

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 16.

Remarks: Nomen nudum, no valid standing.

Nerita leachii

Recluz, M. C. A. 1841. Description de quelques nouvelles

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. leachii, cont.

especies de nerites vivantes. Rev. Zool. Soc. Cuvier.
1841, p. 312.

Description: Testa ovato-oblonga, nitidula mutica nigra,
maculis luteis triangularibus minutis, subserialibus
picta, interdum medio luteo zonata; spira exsertiuscula,
erosa, apertura angusta albida; labio coarctato, piano
compresso, arcuato, longitudinaliter obsolete crenato;
dente cardinali maxima.

Type locality: Nouvelle-Hollande.

Nerita leguillouana [LeGuillouana]

Recluz, M. C. A. 1842. Description de plusieurs especies de
nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842,
p. 178.

Description: T. ventricosa-globosa, crassa, lutea, nigro-
undata, sulcato-costata, longitudinaliter oblique et tenue
striata; spira prominenti, conico-acuta; apertura rotun-
data, superne dilatata, fauce flavida; labio piano, rugoso
et granulos, margine 3-4 dentato; labio crenulato, intus
sulcato, versus spiram bidentato.

Type locality: "Malaisie et les iles Salomon."

Nerita lessonii

Recluz, M. C. A. 1841. Description de quelques nouvelles
especies de nerites vivantes. Rev. Zool. Soc. Cuvier.
1841, p. 311.

Description: Testa globosa, castanea, minima; subepidermide
maculis hastatis, signata; anfractu unico, supra medium
anguloso, spina brevissima instructo; apice oculato;
pertremate rotundato; columella semilunari, margine
arcuata dentibus tribus exiguis notata; suprema maxima.

Type locality: Sicile.

Nerita leucozonias

Gmelin, J. F. 1791. Systema naturae, 13th edition,
p. 3672.

Description: N. testa subglobosa solida badia albo
fasciata; spira depresso-
sculpta.

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. leucozonias, cont.

Type locality: Not given.

Nerita liciniaMuller, O. F. 1774. Vermium terrestrium et fluviatilium.
Vol. 2, p. 178.Description: Nerita testa ovata, incarnata, striata;
apertura in puncta admata.

Type locality: Not given.

Nerita ligataMuller, O. F. 1774. Vermium terrestrium et fluviatilium.
Vol. 2, p. 181.Description: Nerita testa ventricosa, acuminata, subumbili-
cata, fasciis rubris, apertura circulari.

Type locality: "In Museo Spengleriano."

Nerita lineataMuller, O. F. 1774. Vermium terrestrium et fluviatilium.
Vol. 2, p. 189.Description: Nerita testa subulata fusca, lineis septum
spiralis impressis, apertura ovata.

Type locality: "In Museo Spengleriano."

Nerita lineata

Meuschen, F. C. 1787. Museum Geversianum, p. 272 and 274.

Remarks: Not acceptable.

Nerita lineataGmelin, J. F. 1791. Systema naturae, 13th edition,
p. 3684.Description: N. testa intus labioque utroque albis, anfrac-
tibus teretibus striis nigris parallelis concisis; inter-
sticiis striarum violaceis, labio exteriore intus striato.

Type locality: "Sreto malaccensi." —

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. lineata, cont.

Remarks: See Cernohorsky for comments.

Nerita lirellata

Rehder, H. A. 1980. The marine mollusks of Easter Island (Isla de Pascua) and Sala y Gomez. Smithsonian Contrib. Zool. 289, p. 23.

Description [Diagnosis]: From the closely related Nerita morio (Sowerby), this species differs in having a dull exterior surface sculptured with regular spiral furrows and ridges, a lower spire, a columellar lip with only one very obscure tooth, and an operculum whose calcareous part is more minutely granulose, and almost uniformly gray rather than orange with blackish margin.

Type locality: Easter Island.

Nerita listeri

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 177.

Description: Testa parva, semiglobosa, striato-costata, nigerrima, maculis niveis parvulis picta; spira promimula, obtusiuscula; apertura exlutescente, labio plano, laevi, margine bidentato; labro margine tenui, crenulato, nigro, zona viridescente instructo, subtus lacteo, incrassato integerrimo.

Type locality: "Hab. rapportee par M. Hanet Clery de l'Ocean pacifique (M. Petit)."

Nerita listeri

*Pfeiffer, L. 1840. Arch. Naturges. 4, p. 255. [not seen]

Nerita litterata

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3685.

Description: N. testa subglososa alba characteribus inscripta; labio interiori crenulato dentato.

Type locality: "Habiata rara in India." —

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita littoralis

Linnaeus, C. 1758. Systema naturae, p. 777.

Description: N. testa laevi, vertice carioso, labris edentulis.

Type locality: "Habitat ad maris Europaei littore scopulosque, modis innumeris variegata."

Remarks: Now Littorina obturata, Vermeij.

Nerita littorea

*Muller, O. F. 1776. Z. Dan. Prod. p. 244. [not seen]

Nerita litorata

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 315.

Description: Testa ventricosa-oblonga, nigro-violaceouscente, lineis tenuissimis, brevissimis, interruptis transversim seriatis picta; anfractibus 4-5 superne angustatis; spira exsertiuscula, apice acuta; columella coarctata, convexa, carneola, in medio tenue arcuata, crenulata; labro transversim oblique semi-ovali.

Type locality: "Les iles Philippines?"

Nerita livida

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita longii

Recluz, M. C. A. 1842. Description de plusieurs especes de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 178.

Description: Testa ventricosa-globosa, costis depresso 26-27 cincta, dilute ravaida, nigro-trifasciata et maculata: maculis subquadratis longituninalibus seriatim undatis; spira prominenti, decorticata; labio rugoso et verrucoso, aestate, compresso-plano et postice rugis elevatis marginato, antice flavicante tridentato; labro intus breviter striato et bidentato, superne vix dilatato.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. longii, cont.

Type locality: "Hab. la cote de Malabar, a Bombay,"

Nerita longispina

Recluz, M. C. A. 1841. Description de quelques nouvelles especies de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 312.

Description: Var. a). Dilatata spinis coronata.

Var. b). Mutica. Dilatata, spinis nullis, rufa; columella, aurantia, medio arcuata, edentula.

Var. c). Mutica. Ovata, spinis nullis, rufa, zonis nigrescentibus transversis picta, columella laevissime angulosa.

Type locality: Not given.

Nerita luctuosa

Recluz, M. C. A. 1841. Description de quelques nouvelles especies de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 317.

Description: Testa ovato-oblonga, lineis undulatis crebris nigris longitudinalibus picta; angractibus quaternis; ad suturam submarginatis; spira exsertiuscula, subacuta; apertura externe ovata-acuta, intus fusco-viridula; columella plano-concaviuscula, angusta, semi-lunari, obsolete crenulata.

Type locality: New Guinea.

Nerita lugubris

*Voig, F. S. 1834. In Cuvier, Thierreich 3, p. 215. [not seen]

Nerita lunulata

Muller, O. F. 1774. Vermium terrestrium et fluviatilium. Vol. 2, p. 180.

Description: Nerita testa cylindracea alba, reticulata; apertura remota.

Type locality: "In Museo Fabriciano." —

TABLE T. (Continued) CATALOG OF RECENT NERITA SPECIESNerita lutea

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita maculata

*Salis, C. U. von. 1793. Reise Neapel, p. 379. [not seen]

Nerita maculata

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 19.

Remarks: Nom. nudum, no valid standing.

Nerita maculata

Pease, W. H. 1867. Descriptions of sixty-five new species of marine gasteropodae, inhabiting Polynesia. Am. J. Conchol., n.s., 3(4), p. 286.

Description: T. globosa, rotundata, solida, crassa, spiraliter tenuiter et creberrime, linearis, sulcata; spira plano-depressa; area columellari vix concava, granulata et corrugata, fortiter dentata; labro intus valde crenato et dentato; nigra, punctis luteis lentiginosa.

Type locality: Tahiti.

Nerita maculifera

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 345.

Description: Testa globoso-acuta; anfractibus quaternis convexis, transversim sulcatis, longitudinaliter stiratis, violaceis, costis maculis intensioribus articulatis; spira exsertiuscula cuta; apertura extus ovato-acuta, intus semi-lunari; labio plano, inerme, obsolete bidentato; labro ad marginem obsolete crenato, intus omnino inerme.

Type locality: Not given.

Nerita maculosa

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita magdalena

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3677.

Description: *N. testa late sulcata nigra intus alba; labio
flavicante, operculo laevi bidentato..*

Type locality: "Habitat rario circa insulas Magdalena,
...."

Nerita malaccensis

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3684.

Description: *N. testa sulcata intus flavicante; vertice
prominulo, labio exteriori inermi exterius crenulato,
interiore flavicante laevi.*

Type locality: "Habitat in freto malaccensi."

Nerita mammilla

Linnaeus, C. 1758. *Systema naturae*, p. 776.

Description: *N. testa umbilicata ovata glabra, umbilico
obtecto, apertura ovata.*

Type locality: Barbados.

Remarks: Now Polinices or Natica; see Cernohorsky, naticids
of Fiji, 1971.

Nerita marginata

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3678.

Description: *N. testa tenui decussatim striata tuberculata
nigra ochroleuco-maculata subglobosa; apertura extrinsecus
marginata.*

Type locality: Not given.

Remarks: See Zoological Record 1877:Moll 45, marginata
synonym of N. polita.

Nerita marmorata

Meuschen, F. C. 1787. *Museum Geversianum*, p. 270.

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. marmorata, cont.

Remarks: Not acceptable.

Nerita marmorata

Reeve, L. A. 1855. *Conchologia iconica: or, illustrations of the shells of molluscous animals* 9 (*Nerita*), plate 16, species 67.

Description: *Ner. testa orbiculari-depressa, crassiuscula, spira obtuse impressa, anfractibus convexis, versus aper-turam latis, irregulariter spiraliter liratis et linear-i-sulcatis, apertura expansa, labro effuso, columella declivi, interdum minute denticulata; caeruleo-nigro sparsim marmorata.*

Type locality: Not given.

Nerita maroccana

*Salis, C. U. von. 1793. *Reise Neapel*, p. 379. [not seen]

Nerita marochiensis

Gmelin, J. F. 1791. *Systema naturae*, 13th edition, p. 3673.

Description: *N. testa subglobosa oblique plicata; spira mucronata, umbilico bifido.*

Type locality: Not given.

Remarks: Now a Natica, still good west African species name.

Nerita martensiana

Pilsbry, H. A. 1901. *New Japanese marine, land and fresh-water Mollusca*. Proc. Acad. Nat. Sci. Philadelphia, p. 385-408.

Description: Shell globose, small, solid, rather bright sulphur yellow, paler and somewhat mottled with gray or blackish toward the aperture. Surface dull, sculptured with low, rather coarse spiral cords, about 15 on the last whorl, the upper one appressed against the preceding whorl. Spire short, whorls about 3, the last a little depressed below the suture, which is bordered below by a somewhat more prominent cord. Aperture semicircular, yellow or whitish; lip-rib smooth, with a small tubercle

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. martensiana, cont.

above, and another well within near the base of the columella. Columellar area white or yellowish, flat and smooth, the outer border well defined; edge of columella straight, with two or three low, subobsolete teeth, the upper one strongest.

Type locality: LooChoo Islands (=Ryukyu Islands).

Remarks: Vermeij, color form of N. helicinoides.

Nerita maura

Recluz, M. C. A. 1842. Description de plusieurs especes de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 179.

Description: Testa ovata, ventricosa, crassa, costis depresso 33-35 cincta, luteo-fluva, maculis transversis longitudinaliter undatim digestis picta; anfractibus superne depresso; spira laterali, brevissima; apertura alba, fauce luteola; labio largo, plano, basi rugoso, antice tridentato; labro extius superneque leviter planulato, intus tenuiter sulcato, edentulo.

Type locality: Hab. la Polynesie, aux iles Saint-Pierre et Saint-Francois (Peron et Lesueurs), a Madagascar (Mus. Paris).

Remarks: Vermeij says this is a valid name for a ribbed maxima type, black.

Nerita mauritiae

Recluz, M. C. A. 1841. Description de quelques nouvelles especies der nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 106.

Description: N. testa semiglobosa, crassiuscula, nigerrima, transversim tenue striata, costis latis planis, maculis albis remotis impressis insculpta; anfractu infimo superne depresso; spira priminula, rotundato-obtusa, eroso-albida; apertura exalbidi, rotundata, labio angusto, bidentato, supra rugoso et verrucosa; labro crasso, intus striato, superne vix dentato.

Type locality: Les iles Maurice et Rodriguez.

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESNerita maxima

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3683.

Description: *N. testa solida crassa glaberrima radiis
undulatis atris et flavicantibus picta; labio exteriore
edentulo, interiore subconcavo quadridentato.*

Type locality: Unknown.

Nerita melanoleucas

Meuschen, F. C. 1787. *Museum Geversianum*, p. 268.

Remarks: Not acceptable.

Nerita melanostoma

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3674.

Description: *N. testa tenui pellucida laevi oblonga; an-
fractu primo ventricoso plano maximo reliquis compressis,
umbilico semiclauso.*

Type Locality: "Habitat in mari indico," p. 3674.

Remarks: Possibly Polinices (Mammila).

Nerita melanotragus

Smith, E. A. 1884. *Mollusca*, p. 69. In Report on the
zoological collections made in the Indo-Pacific ocean
during the voyage of H.M.S. "Alert" 1881-2. London.

Description/Remarks: Based on a description of N. atrata of
Reeve, 1855; nothing added.

Nerita menkeana

Recluz, M. C. A. 1842. Description de plusieurs especes de
nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842,
p. 183.

Description: *T. ovata, semiglobosa, ventricosa, striata,
olivaceo-nigricante, maculis parvulis flavescentibus dense
adspersa; anfractibus 3-4; ultimo depresso et anguloso; ad
angulum spinis rarissimis brevibus notato; spira prominula,
convexo depressa, decorticata; apertura intus carneolata,
marginis flava; labio lutescente, convexo-depresso, antice*

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. menkeana, cont.

recto, dente cardinali tantum signato.

Type locality: Tahiti.

Remarks: Described in subgenus Clithon.Nerita mertoniana

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 71.

Description: Ner. testa ovato-globosa, maculis pallide lutescentibus zonisque articulatis nigro-purpurascensentibus fimbriatis cincta; spira vix prominente, rotundata, apice eroso; apertura obliqua, luteola; labio subconvexo, angusto, margine tenue arcuato et crenulato.

Type locality: "Hab. cum praecedente."

Remarks: Affinities to Nerita ovalaniensis Lesson; probably fresh or brackish water species.

Nerita michaudi

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 315.

Description: Testa subglobosa, tenui, olivaceo-fuscescente, lineolis nigrescentibus angulato-flexuosis picta; spira erosa, obliquissima; apertura rotundata, albido-caerule-scente; labio angusto, planulato, in medio vix arcuato et crenulato; labro superne subarcuato, anterius subtruncato.

Type locality: Unknown.

Remarks: See also Recluz (1843, p. 72) for mention of N. michaudiana.

Nerita miliacea

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p 316.

Description: Testa minutissima, tenuissima, hyalina, lucida, subglobosa, transversim lineis binis aurantiis fasciata, interstitiis lineolis purpureis undatis notata, superne fasciculis purpureis radiata; spira vix prominula,

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. miliacea, cont.

obtusata; labio convexiusculo, medio crenato.

Type locality: Not given.

Nerita millepunctata

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 19.

Remarks: Nomen nudum, no valid standing.

Nerita minima

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 346.

Description: Testa minima subglobosa; dilute olivacea, sphacaelis nigro-marginatis undatusque picta; spira punctiformi apertura dilatata, rotundata, inuts semi-lunari; labio angusto, plano, obsolete crenulato.

Type locality: Nukuhiva.

Nerita minuta

Muller, O. F. 1774. Vermium terrestrium et fluviatilium, Vol. 2, p. 179.

Description: Testa subvirescens, cylindrica, aequali sere crassitie, vertice obtuso; anfractus tres crum semisse, teretes; major reliquis duplo longior; apertura ovalis, margine aequali.

Type locality: Not given.

Nerita minuta

Sowerby, J. D. C. 1825. The mineral conchology of Great Britain; or , Vol. 5, p. 93.

Description: Hemispherical, smooth; spire indistinct; aperture oval; columellar lip not toothed.

Type locality: "Figures from the Cabinet of the Rev'd George Cookson" (p. 93).

Nerita mitchelli

Oldroyd, I. S. 1933. Two interesting shells from the Philippine Islands. Philippine J. Sci. 52(2), p. 205.

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. mitchelli, cont.

Description: Shell thick, with fifteen to eighteen fine spiral riblets, the interstices of equal width. Exterior dirty white, banded with dull rose; in some cases the color nearly covers the shell while in others only part of the body whorl is rose. Aperture porcellaneous and polished, outer lip thickened with faint dentations within. Columellar lip with from three to four dentations, the type having three; the columellar area with fine ridges covering the entire area and highly polished.

Type locality: Philippine Islands, no definite locality label.

Remarks: Type--No. 16368, Bureau of Science; Paratype at Conchological Collection, Leland Stanford Junior University.

Nerita mittreana

Recluz, M. C. A 1842. Description de plusieurs especes de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 181.

Description: T. ovato-rotundata, ventricusa, tenui, glabruscula, nitida, cornea, maculis oblongis, albis, alternantibus et interstitiis fusco-caeruleis longitudinaliter lineolatis picta; spira prominula, latiuscula, subcentrali, planulta; apertura flavescente; labio piano, edentulo, margine recto, acuto; labro superne et inferne saepius nigro marginato.

Type locality: "Hab. les eaux douces de Grasse en Provence"

Remarks: Described in subgenus Neritina.

Nerita monile

Bolten, J. F. 1798. Musuem Boltenianum, Part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita montacuti

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 174.

Description: N. testa ventricosa-globosa, supra medium

TABLE T. (Continued) CATALOG OF RECENT NERITA SPECIESN. montacuti, cont.

spinis brevibus coronata, oblique striata, lutescente, superne et infra medium zonis nigris lutescente maculatis picta; suprema latiore; anfractibus tribus superne depresso-planis; spira depresso-convexa; apertura dilatata, rotundata, albido-lutescente; labio angusto, medio concavo, margine denticulato, in centro arcuato.

Type locality: Island of Burias, on stone in a rivulet.

Nerita moreleti

Recluz, M. C. A. 1845. Description de mollusques terrestres et fluviatiles du Portugal. Rev. Zool. Soc. Cuvier. 1845, p. 311.

Description/Remarks: No description; renaming of Nerita violacea Gmelin; freshwater species.

Nerita multiungis

Menke, K. T. 1847. Verzeichniss einer sendung von chonchyliden von Mazatlan, mit einigen kritischen bemerkungen. Z. Malakozool. 4, p. 179.

Description: N. testa subglobosa, solida, nigra, cingulis 32 angustis circumvallata; spira prominula; labio convexo, quasridentato, rugoso, dentibus intermediis distinctis obtusiusculis, interstitiis dentium rugarumque flavescentibus.

Type locality: Not given.

Nerita muricata

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 20.

Remarks: Nomen nudum, no valid standing.

Nerita musica

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 19.

Remarks: Nomen nudum, no valid standing.

Nerita musiva

Gould, A. A. 1848. [untitled]. Proc. Boston Soc. Nat. Hist. 2, p. 238.

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. musiva, cont.

Description: Testa modica, subsolida, transversa, ovoidea, ventricosa, fusco-cinerea, longitudinaliter crasse striata, sulcis volventibus ordinatis arata, liris interpositis planulatis, sub-granulosis, lineolis crebris albidis sectis; apertura sub-circularis, caliciformis, peristomate sub-continuo; labor edentulo, albo, sulcis obliquis fuscis tenuiter arato; labio concavo, fusco, granulis paucis notato, denticulis tribus instructo.

Type locality: Collected during the U.S. Exploring Expedition.

Remarks: Bishop Museum copy marked "=signata, (Macleay)."

Nerita mustelina

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita navicellina

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 346.

Description: Testa suborbiculari, posterius truncata, convexo depressa, tenuissima sublente transversim tenue sulcata et longitudinaliter obsolete striata; vertice minimo, obliquo, marginali; labio piano, margine tenue excavato et obsolete crenulato; labro superne dilatato, subaruiculato.

Type locality: "Hab. Hamoa" (Samoa?).

Nerita nebulata

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 71.

Description: Ner. testa ovato-globosa, tenuiter striata, luteo-fuscescente, lineis nigris squamaeformibus parvulis nebulata; spira prominula, rotundata; labio angustato, subconvexo, margine retiusculo, crenulato, flavescente; labro semi-ovato, intus caerulescente, margine flavo.

Type locality: Immimaylan, in a mountain stream.

Remarks: Freshwater species.

TABLE T. (Continued) CATALOG OF RECENT NERITA SPECIESNerita nebulosa

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita neritinaoides

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 5, species 21.

Description: Ner. testa ovata, subangusta, solida, spira leviter obtuse exserta, anfractibus superne concavo-declivibus, laevigatis, apertura simplici, area columellari concavo-planata, obsolete minure denticulata; intense nitente-atra, area columellari pallide aurantia.

Type locality: Not given.

Nerita neritopsoidea

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 16, species 69.

Description: Ner. testa ovato-turbinata, spira exserta, mucronata, anfractibus rotundatis, conspicue dense spirali-ter liratis, labro intus incrassato, minute crenulato, dente minute laterali, columella tumidiuscula, oblique corrugata, fortiter dentata; pallide straminea.

Type locality: Not given.

Nerita nigella

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

Nerita nigreola

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 20.

Remarks: Nomen nudum, no valid standing.

Nerita nigerrima

Dillwyn, L. W. 1817. A descriptive catalogue of recent shells arranged according to the Linnaean method with particular attention to the synonymy, p. 996.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. nigerrima, cont.

Description: Shells sub-globular, smooth, thick, opaque, and minutely striated transversely; outer lip entire, and the inner lip slightly wrinkled.

Type locality: Not given.

Remarks: Species from Chemnitz, p. 309.

Nerita nigrescens

Colbeau, J.-A.-J. 1859. Materiaux pour la fauna malacologique de Belgique 1, p. 11.

Description: Not given.

Type locality: Not given.

Remarks: Described as Nerita fluviatilis, L. variete nigrescens; European Theodoxus.

Nerita nigrita

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 19.

Remarks: Nomen nudum, no valid standing.

Nerita nigrocincta

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita nitida

*Donovan, E. 1803. British shells, pl. cxliv. [not seen]

Nerita nivea

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita nivosa

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 16, species 66.

Description: Ner. testa ovata, spira subexserta,

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESN. nivosa, cont.

anfractibus juxta spiram declivibus, deinde convexis,
spiraliter impressis et linearis-sulcatis, sulcis valde
irregularibus, labro intus subincrassato, dente unico
lateralis, columella declivi, obsolete corrugata, dentata;
intese caeruleo-viridi, albo sparsim nebulata.

Type locality: Not given.

Nerita nodosa

Linnaeus, C. 1758. Systema naturae, p. 777.

Description: N. testa nodosa pluribus ordinibus soldia,
labro angulato.

Type locality: Not given.

Nerita notata

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita novae-guineae

Lesson, R. P. 1830 [1831]. Zoology, Vol. 2, p. 371. In
Voyage autour du monde, ..., La Coquille, ...

Description: Cette petite coquille a 7 lignes de hauteur
sur 8 de diametre; son test est epais, dilate au-dessus
de la levre droite, a spire conique, saillante. Elle est
couverte de cotes arrondies, rapprochees, tres-lisses,
et separees par un sillon au milieu duquel regne une
ligne cordonnee de meme forme, mais plus petite que les
cotes qui l'elabore. Sa levre droite est en demi-cercle,
mince et lisse a son bord, puis dilatee et garnie de
petites aretes regulieres et de sillons jusqu'en haut,
ou s'elevent deux dents. La columelle est courte, droite,
ridee, ayant huit cotes se terminant par quatre fortes
dents.

Type locality: Dorery, New Guinea.

Nerita novae-hiberniae

Lesson, R. P. 1830 [1831]. Zoology, Vol. 2, p. 371. In
Voyage autour du monde, ..., La Coquille, ...

Description: Cette nerite a l'aspect de la Nerita

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. novae-hiberniae, cont.

malaccensis, mais elle s'en distingue par plusieurs caracteres. Sa spire est conique et saillante, ce qui n'a pas lieu dans la malaccensis. Sa levredroite est beaucoup plus mince, plus concave, et son rebord est tresbas. Une seule dent marque la haut de cette levre qui est lisse et n'a qu'un ressaut circulaire interne descrivant comme elle un demi-cercle. La columelle est plate et nue, marquee seulement d'une a trois echancrures et dents peu nettes. Sa surface est courverte de cotes saillantes, arrondies, regulieres, separees par un sillon creux et lisee. Aucune strie longitudinale n'apparait en travers sur les cotes.

Type locality: New Ireland [Praslin, a la Nouvelle-Irlande].

Nerita novae-caledonieNerita nubecula

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

Nerita numidica

Recluz, M. C. A. 1841. Description de quelques especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 342.

Description: Testa semiglogosa ennuta, tenui, nigrata seu vinacea, nitida, punctis minimis, pallidia obsolete notata seu seriatim tranverse fasciata; spira laterali, saepius erosa; anfractibus superne rotundatis, apertura orbiculari, subcaerulecente, columella angusta, plana medio subarcuata, edentula.

Type locality: "Hab. Oran deans l'Atlas, vallee de Tisi (Tervier)."

Nerita nuttalli

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 276.

Description: Testa dilatata, suborbiculari, ventricosa, transversim tenue striata, nigerrima, maculis albido-fulvis triangularibus varia; spira retusa, eroso albido; labio plano, vix arcuato, crenulato; labro intus albido-caerulecente, auriculato: auricula dextra convexo angulosa, posterius oblique truncata.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. nuttalli, cont.

Type locality: "Les îles Sandwich (MM. Petit et Janelle)."

Nerita nux-castanea

Martyn, T. 1789. Universal conchologist 3, fig. 106.

Remarks: Invalid, rejected name.

Nerita obatra

Recluz, M. C. A. 1851. Description de quelques coquilles nouvelles. J. Conchyliol. 2, p. 203.

Description: N. testa transversim oblonga, ventricosa, striis decurrentibus argute impessa, nigra, maculis obscurisa variegata; anfractu infimo superne depresso plano; spira extersiuscula, conico acuta; apertura ad peripheriam lutescente; labio calloso, plano laevi, margine tridentato; labro intus breve crenulato.

Type locality: "Hab. Nous l'avons trouvée avec notre Nerita essingtoni var. totaroseo-rubra, et autres espèces venant des côtes de Taiti."

Nerita obscurata

Recluz, M. C. A. 1842. Description de plusieurs espèces de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 183.

Description: T. ventricoso-globoso, longitudinaliter oblique et dense rugosa, subuplicata, fusco-nigra, maculis parvulis lutescentibus sub oculo et lumine notata; anfractibus 2 1/2; ultimo spinis nigris brevibus ostendente; spira retusa, partim decorticata; labio flavescente, superne calloso, antice in medio arcuato et ruguloso.

Type locality: Tahiti.

Remarks: Described in subgenus Clithon.

Nerita ocellata

LeGuillou M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1842, p. 344.

Description: Testa ventricoso-globosa, substriata; nigra, maculis transversim oblongis pallide croce ispicta; spira erosa, apertura ovato-rotundata, columella plana posterius

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. ocellata, cont.

erosa, medio vix granulosa, margine sub-bidentata, labro acuto, intus sulcato, superne tenue bidentato.

Type locality: Sandal-bay [?].

Remarks: Vermeij--that name still may used by some.

Nerita oculata

Meuschen, F. C. 1787. Museum Geversianum, p. 274.

Remarks: Not acceptable.

Nerita oculus

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita oleagina

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 10, species 47.

Description: Ner. testa ovato-globosa, spira exserta, mucronata, anfractibus convexis, spiraliter elevato-striatis, labro intus incrassato, vix crenulato, dente unico laterali, columella convexiuscula, promiscue verrucoso-corrugata, fortiter dentata; nigrante-olivacea, interdum nigro-fasciata, maculis albidis undatis nigro-umbratis oblique veriegata.

Type locality: Not given.

Nerita olivacea

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 17.

Remarks: Nomen nudum, no valid standing.

Nerita olivacea

Recluz, M. C. A. 1842. [untitled]. Proc. Zool. Soc. London 10, p. 172.

Description: N. testa ovato-globosa, ventricosa, crassa, griseo-olivacea, anfractibus 2 1/2 superne depresso-lis; apice subconvexo, obtuso, laterali, in medio deroso;

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. olivacea, cont.

apertura extus subquadrata, intus exalbida, labio postice rugescente, plano, margine crenato et in medio vix arcuato.

Type locality: Agoo, province of Pangasinan, island of Luzon; on stones in a rapid river.

Nerita olivaria

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 344.

Description: Testa ventricoso-globosa, superne depressiuscula, cinereo-olivacea, maculis albis nigro marginatis transversim obsolete fasciata; spira brevissima; apertura dilute straminea; labio plano, inerme, margine in medio dentibus 2, 3, 4 obsoletis, exiguis; labro intus tenui sulcato et unidentato.

Type locality: Mindanao.

Remarks: See Mienis (1970, p. 63) for designation of N. junghuni Shepman, 1901, lectotype.

Nerita opaca

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

Nerita orbignyana

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 108.

Description: Testa subovata, ventricosa, superne et inferne coarctata, pulchre picta, nitida, longitudinaliter subtilissime striata; spira brevissima, rotundata, subactua; labio angusto, plnao, declivi, in medio obtuse obsoleteque bidentata, labio extus et intus integerrimo.

Type locality: La mer Rouge, dans le canal de l'Yemen.

Remarks: Vermeij says valid Red Sea subspecies of N. polita; also N. orbignyana n. sp. as fossil from Cotteau (1854).

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita orientalis

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3673.

Description: *N. testa subglobosa nitida laevissima; spira
base subrugosa, columella nivea.*

Type locality: "Maribus orientalibus."

Remarks: See Cernohorsky.

Nerita ornata

Sowerby, G. 1823. *The genera of recent and fossil
shells, Nerita*, fig. 4.

Description: *Testa subglobosa, transversim sulcata, albo
vel luteo-nigroque variegata, spira prominula; labio
interno quadridentata, superne rugoso.*

Type locality: Not given.

Remarks: Sowerby thought this species to be *N. histrio*
Linnaeus but because descriptions so variable and figures
so bad, felt it impossible to determine; see also Wood
(1828); now may be junior synonymy of *scabracosta* of Gulf
of California or species from west coast of Panama.

Nerita oryzarium

Recluz, M. C. A. 1841. *Description de quelques nouvelles
especes de nerites vivantes. Rev. Zool. Soc. Cuvier.*
1841, p. 178.

Description: *Testa semiglobosa, seu ovato-subglogosa,
solida, laevi, lineis obsoletis remotis transversis in-
sculpta; olivacea, maculis albis seu ravidis flammulata
variegataque; spira spominula, subactua; labio compresso-
plano, granulato, posterius rugoso, marigne 2,3,5 dentato;
labio integro, intus striato, superne bidentato; dente
secunda saepe irregulari.*

Type locality: Bombay.

Remarks: Valid Indian species; see Reeve (1855, pl. 10, sp.
46) for figure.

Nerita otaitensis [o-taitensis]

Lesson, R. P. 1830 [1831]. *Zoology*, Vol. 2, p. 370.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. otaitensis [o-taitensis], cont.

Description: Cette nerite est epaisse; solide, arrondie, a spire courte, a dernier tour gaarni de cotes regulieres, separees par un sillon creux. Bouche etroite, a levre externe lisse interieurement, arrondie, garnie en bas de cinq dents coniques, egales. Bord columellaire droit, marque de quatre plis, gros, lisses, deparés par un sillon profond.

Type locality: Tahiti and BoraBora.

Remarks: Vermeij--synonym of N. plicata.

Nerita owenii

Wood, W. 1828. Index testaceologicus, Supplement, p. 26.

Description: Not given.

Type locality: Africa.

Nerita oweniana

Wood, W. 1828. Index testaceologicus, Supplement, p. 25.

Description: None given.

Type locality: Africa.

Nerita panayana

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 170.

Description: N. testa parva, ovata-hemisphaerica, ventricosa, postice angustata, tenui; luteo-viridescente, lineolis nigrescentibus angulato-flexuosis creberrimis picta; anfractibus 1 1/2 supremo eroso-retuso; apertura extus orbiculari, intus dilatata, albido-caerulecente; columella semi-lunari plana, albo, antice rectiuscula.

Type locality: "River Dingle, province of Ylo-Ylo [Iloilo], island of Panay; on small stones."

Nerita panvensis

Recluz, M. C. A. 1844. [untitled]. Proc. Zool. Soc. London 11, p. 200.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. panayensis, cont.

Description: Ner. testa parva, ovato-conica, aurantia, transversim sulcata, longitudinaliter crebre striata; costis subter lente subgranosis; spira conico-acuta; labio plano, lacteo, dentibus tribus remotis instructo; labro margine crenao, intus calloso, lacteo, plane laevissima.

Type locality: "Ilo-Ilo, province of Panay; under stones at low water" H. Cuming.

Nerita papilio

Bolten, J. F. 1798. Museum Boltenianum, part 2, p. 19.

Remarks: Nomen nudum, no valid standing.

Nerita papilla

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3675.

Description: N. testa pelludica tenui oblonga decussatim striata sordide lutea; apertura subovali, columella alba, umbilico demicauso.

Type locality: "Habitat frequens ad Tranquebariae littora, anfractibus 4."

Nerita papuana

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 105.

Description: N. testa ovato-subglobosa, nigerrima, transversim striato-costata; costis inaequalibus, geminatis; anfractu ultimo superne depresso; spira prominula, apice subacuta, albida; apertura rotundata, alba; labio rugoso, margine tridentato, dente prima maxima; labro intus striato, superne bidentato.

Type locality: New Guinea.

Nerita paralella

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 16.

Remarks: Nomen nudum, no valid standing.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita parvula

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 346.

Description: Testa parvula, subglobosa, laevissima, nitula, nigerrima; spira convexa, apice eroso; apertura subdilata; labio angusto, plano, margine superiore obsolete crenato; operculo olivaceo, laevi.

Type locality: Lebouka (Viti) [=Levuku, Fiji?].

Nerita patella

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 16.

Remarks: Nomen nudum, no valid standing.

Nerita patula

Wood, W. 1828. Index testaceologicus, Supplement, p. 25.

Description: None.

Type locality: Not given.

Nerita patula

Recluz, M. C. A. 1841. Descriptions de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 148.

Description: Testa ovato-subglobosa, anterius dilatata, tenui, transversim striat; striis angustis granosis, grisea, albo, nigro, et ravid variegata seu fasciata; spira brevissima obtusa; apice flavescente; apertura patula, dilute aurantia; labio lato, subconcavo, nigro maculato, granulato, margine arcuato, subedentulo; labro dilatato, intus striato, edentulo.

Type locality: Not given.

Remarks: Apparently valid species from Indonesia and Philippine Islands, looks like N. chameleon.

Nerita pecten

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 20.

Remarks: Nomen nudum, no valid standing.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita pellis-armeniana

Martyn, T. 1789. Universal conchologist, vol. 3, p. 109.

Remarks: Nonbinomial, invalid, rejected name.

Nerita pellis-serpentis

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 19, species 83.

Description: Ner. testa ovato-turbinata, spira exserta, mucronata, anfractibus rotundatis, undique desne liratis, apertura parva, columella obsucra dentata; albida, punctis rufis dense promiscue tessellata.

Type locality: Not given.

Nerita pellucida

Adams, J. 1797. The specific characters of some minute shells discovered on the coast of Pembrokshire, with an account of a new animal. Trans. Linnean Soc. 3:67.

Description: N. testa laevi pellucida, anfractibus tribus.

Type locality: Pembrokshire?

Nerita peloronta

Linnaeus, C. 1758. Systema naturae, p. 778.

Description: N. testa striata, labiis dentatis: interiore planiusculo rugoso.

Type locality: "Habitat in O. Asiae ad Bandam."

Remarks: See Russell (1941), OK in west Atlantic.

Nerita pennata

*Born, J. von. 1778. Index Mus. Caes. Vind, p. 420. [not seen]

Nerita pennata

*Davis, H. 1791. In T. Pennant. Ind. Zool., 2 ed., p. 158. [not seen]

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita perrottetiana

Recluz, M. C. A. 1841. Description de quelques de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 337.

Description: Testa ovato-semiglobosa, fusco-nigricante, tenuissime striata; anfractibus tribus; ultimo superne coarctato; spira brevi, convexa, obtusata; apice aplido, sub-hyalino; apertura extus rotundata; labio plano, albido, in medio vix arcuato obsoleteque crenato.

Type locality: "Hab. Dans les rivieres des monts Neelgheries,"

Nerita peruviana

Philippi, R. A. 1844. Abbildungen und beschreibungen neuer oder wenig gekannter conchylien, unter mithulfe mehrere Deutscher Conchyliologen, herausgegeben von..., p. 84.

Description: N. testa ovata, laevi, opaca; ex albo et atro variegata; spira prominente; anfractu ultimo demum ad suturam depresso; labro 9-10 dentato; columella 4-dentata, dentibus medianis minutis; labio laeviusculo, concavo, granulis obsoletis notato.

Type locality: Peru, legit frater E. G. Philippi.

Nerita perversa

Linnaeus, C. 1758. Systema naturae, p. 3686.

Description: N. testa cingulata: spira contraria, apertura octodentata.

Type locality: Not given.

Nerita petechialis

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

Nerita petitii

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 273.

Description: Testa ovato-subhemispherical, magna, ventricosa, laevigata, viridi-fuscescente; spirae anfractibus

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. petitii, cont.

binis, vix exsertis, rotundato-obtusis, sutura linearis
distinctis; apertura maxima, superne et inferne dilatata,
externe ovata, intus cinnamomea; columella lata, plana,
margin'e subarcuata ac crenulata.

Type locality: Hab. Saint-Domingue.

Remarks: Neritina?

Nerita pfeifferiana

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc.
London 11, p. 72.

Description: Ner. testa ventricoso-ovata, nigra, albido-lutescente bizonata; zona mediana spiram decurrente; anfractibus quaternis, convexis, ad suturam minime marginatis; apice decorticato, obtusiusculo; apertura rotundata; labio compresso, concaviusculo, albido, in margine arcuatim ac obtuse denticulato; denticulis obsoletis, cardinali productiusculo; labro dilatato, tenui nigro-caerulecente et luteo quadrifasciato.

Type locality: "from New Ireland, in a mountain-stream, by R. B. Hinds, Esq."

Nerita phasiana

Recluz, M. C. A. 1842. Description de neuf especes de nerites nouvelles, suivie d'observations sur les Nerita cornea et dubia. Rev. Zool. Soc. Cuvier. 1842, p. 75.

Description: Testa subglobosa, nitida, tenue striata, lineolis nigris creberimis undulatis et maculis lutescentibus picta; spira vix prominula, convexo-depressa, subacuta; sutura profunda; apertura lutescente obliqua; labio calloso, convexo, margine recto et ruguloso; labro basi depresso, superne producto.

Type locality: Saint-Domingue.

Nerita philippii

Recluz, M. C. A. 1841. Description de quelques especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 341.

Description: Testa minuta, semiglobosa, hyalina, nitida, rubro-violacecente reticulata; anfractibus binis seu 2 1/2, supremis albidis, convexo-obtusis, apertura

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. philippii, cont.

rotundata, labio angusto, planiusculo, edentulo.

Type locality: Sicily.

Nerita phrygia

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita phyrgonia

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita pica

Meuschen, F. C. 1787. Museum Geversianum, p. 266.

Remarks: Not acceptable.

Nerita pica

Gould, A. A. 1859. [untitled]. Proc. Boston Soc. Nat. Hist. 7, p. 43.

Description: T. parva, tenuis, oblique ovato-globosa, rudis anthracina maculis parvis multiformibus albis ornata. Spira haud elevata. Apertura semi-circularis, labro simplici nigro marginata; columella excavata, nitida, flavescente, edentato. Operculum virescens, granulatum.

Type locality: Simoda.

Nerita pica

*Lischke, C. E. 1869. Japanische Meeres-conchylien, p. 85.
[not seen]

Nerita picea

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 151.

Description: Test ovata, nigerrima, nitida, transversim striata, anfractu infimo subsuturam depresso, spira brevi, apice acuto seu eroso; apertura subovata, alba; labio

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIES

N. picea, cont.

plano, laevi margine bidentato; labro margine creulato, superne compresso, intus tenue striato, unidentato; dente fissa.

Type locality: Les iles Sandwich.

Nerita pictata

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

Nerita piperina

Dillwyn, L. W. 1819. A descriptive catalogue of recent shells, arranged according to the Linnaean method;..., p. 995.

Description: Shell sub-globular, smooth, thin, of a dull yellowish colour, with triangular black spots, and the inner lip denticulated.

Type locality: Coasts of Malabar (Chennitz).

Nerita piscinalis

Muller, O. F. 1774. Vermium terrestrium et fluviatilium. Vol 2, p. 172.

Description: Nerita testa globosa, perforata, rufo-fusca.

Type locality: Not given.

Nerita pisiformis

Recluz, M. C. A. 1842. Description de plusieurs especes de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 181.

Description: T. parva, subglobosa, nigerrima, interdum maculis oblongis sparsis albis picta, aetate partim corrosula; spira brevi, rotundata; apertura extus sub-rotundata; lbio angusto, semilunari, plano, lutescente, margine vix arcuato, obsolete denticulato; cardinali producto, majori; labro superne subadscendente.

Type locality: Polynesia? O-Tahiti?

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita planospira

Anton, H. E. 1839. Verzeichniss der Conchylien..., p. 30.

Description: Planospira mihi, oval, seitlich zusammengedrückt; Wirbel sehr flach; mit schmalen Rippen: rotlich, mit schwarzen Zic Zac Streifen; Mundsaum scharf; Kanal ziemlich tief; innere Lippe auf der untern Halftie gekornt, auf der oberen gerunzelt, ziemlich stark gezähnelt.

Type locality: Not given.

Nerita plexa

Dillwyn, L. W. 1817. A descriptive catalogue of recent shells, arranged according to the Linnaean method;..., p. 998.

Description: Shell roundish, with transverse crenulated ribs which are alternately larger; inner lip plaited above, and tuberculated below.

Type locality: Coasts of Tranquebar, Ceylon and Nicobar Islands.

Remarks: Vermeij says this is synonym of N. exuvia.

Nerita plicata

Linnaeus, C. 1758. Systema naturae, p. 779.

Description: N. testa sulcata, labiis dentatis; interiore rotundato; exteriore utrinque dentibus acutis conicis.

Type locality: Not given.

Nerita plumbea

Meuschen, F. C. 1787. Museum Geversianum, p. 274.

Remarks: Not acceptable.

Nerita plumbea

Wood, W. 1828. Index testaceologicus, Supplement, p. 25.

Description: None given.

Type locality: Not given.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita politaLinnaeus, C. 1758. *Systema naturae*, p. 778.Description: *N. testa laevi, labiis utrisque dentatis.*

Type locality: "O Asiae."

Nerita pomahakaensisFinlay, H. J. 1924. Some necessary changes in names of New Zealand Mollusca. *Proc. Malacol. Soc. London* 16(2):99-107.Remarks: Nomen novum; *N. nitida* (Hutt.), preoccupied by Donovon's *N. nitida*.Nerita powisianaRecluz, M. C. A. 1843. [untitled]. *Proc. Zool. Soc. London*, 11, p. 71.Description: *Ner. testa ovato-transversa, ventricosa, superne depresso-planiuscula, varie picta; spira brevisima, derosa, auricula elevata, marginata; apertura ovata, dilatata, externe subrecta; labio planissimo, antice recto, acuto et sub lente ruguloso.*

Type locality: New Ireland, in mountain streams, by R. B. Hinds, Esq., surgeon of H.M.S. Sulphur.

Nerita praecognitaAdams, C. B. 1848. *Specierum novarum conchyliorum, in Jamaica repertorum, synopsis*. *Proc. Boston Soc. Nat. Hist.* 2, p. 7.Description: *N. t. crassa, olicacea, albo-marmorata; anf. 3, superioribus convexiusculis; ultimo costulis rotundatis crebris, striatis, numerum (a 20 ad 25) et latitudinem in diversis exemplis disparibus--ornato; labro acuto, a costulis eleganter pectinato, intus juxt marginem polito, interius crasso et crenulato, et supra infraque bituberculato; labio medio excavato et bidentato, alibi plicato vel noduloso. Operculo pallide murino, creberrime papilloso.*

Type locality: Jamaica?

Remarks: Reeve (1855, plate 7, species 30) provided a description of *N. praecognita* and cited C.B. Adams, MS. Mus. Cuming.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita prasina

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita prolifera

*Muller, O. F. 1784. Zool. Dan. 2, p. 34.

Remarks: Nereis?Nerita promontorii

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3686.

Description: N. testa atra ovato acuminata; suclis triginta.

Type locality: "Habitat rario ad littus Africæ inter promontoria Manuel et viride,..."

Nerita proxima

Sowerby, G. B. 1903. Descriptions of fourteen new species of marine Mollusca from Japan. Ann. Mag. Nat. Hist., Ser. 7, 7, p. 498.

Description: Testa oblique ovata, sulva, nigro-fusco variegata, spira parva, abbreviato concia, convexa; anfractus 4-convexi, oblique subtiliter striati, spiraliter valde lirati, liris numerosis, rotundatis; anfractus ultimus latus, obliquus, supra leviter concavus, deinde convexo rotundatus, liris circiter 25 inaequalibus, rotundatis instructus; apertura patula eximie plicata, luteo tincta; labrum acutum, crenulatum, margine dextro oblique descendens; area columellari albo callosa, nitens, leviter irregulariter rugosa, ad marginem tridentata.

Type locality: LooChoo.

Nerita pulchella

Wood, W. 1828. Index testaceologicus, Supplement, p. 26.

Description: None given.

Type locality: West Indies.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita pulchella

Recluz, M. C. A. 1843. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 175.

Description: N. testa subglobosa, nitida, vix striata, luteo-viridescente, maculis nigris sagittatis et zonis aequalibus remotis nigris lutescente articulatis ornata; spira nulla; apice convexo; apertura semi-rotunda, caerulescente; labio angusto, subconvexo, superne calloso, antice subrecto, denticulato; dente cardinali majori.

Type locality: Sual, province of Pangasinan, isle of Luzon; in a small stream.

Remarks: Clithon?

Nerita pulchra

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 19.

Remarks: Nomen nudum, no valid standing.

Nerita pullata

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

Nerita pulligera

Linnaeus, C. 1767. Systema naturae, 12th edition, vol. 1, pt. 2, p. 1253.

Description: N. testa laevi rudi, spirula excavato-oculata, labio interiore laevi crenulato.

Type locality: "Indiae fluviis."

Nerita punctata

Muller, O. F. 1774. Vermium terrestrium et fluviatilium. Vol. 2, p. 190.

Description: Nerita testa subulata, lutescente, fascia juxtasuturam candida striis rubris; anfractibus minoribus sulcatis.

Type locality: None given.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita punctata

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita punctata

Dillwyn, L. W. 1817. A descriptive catalogue of recent shells, arranged according to the Linnaean method;..., p. 981.

Description: Shell sub-globular, whitish, with numerous scattered yellowish dots, and three transverse rows of reddish irregular spots.

Type locality: "Inhabits the Mediterranean."

Remarks: Chemnitz species, p. 173.

Nerita punctata

Quoy, J. R. C., and J. P. Gaimard. 1834. Voyage de decouvertes de l'Astrolabe..., p. 185.

Description: Nerita, testa crassa transversim striata, nigra, punctis albis minimis notata; spira convexa, prominenti; apertura alba, antice denticulata; labio dentato.

Type locality: "Habite i'Ile-de-France, sur le rivage, vis-a-vis l'habitation de M. Charles Talfair, au Mapou."

Nerita punctulata

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita pupa

Linnaeus, C. 1767. Systema naturae, 12th edition, vol. 1, pt. 2, p. 1253.

Description: N. testa laevi subrotunda lactae, anfractibus striis transversis nigris parallelis.

Type locality: Not given.

Remarks: Now Puperita pupa.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita purpurata

Meuschen, F. C. 1787. Museum Geversianum, p. 170.

Remarks: Not acceptable.

Nerita pusilla

Muller, O. F. 1774. Vermium terrestrium et fluviatilium.
Vol. 2, p. 171.

Description: Nerita testa globosa, virescente, perforata,
apertura circinnata.

Type locality: Not given.

Nerita pyrenaica

Moquin-Tandon, A. 1855. Histoire naturelle des mollusques
terrestres et fluviatiles de France, vol. 2, p. 249.

Description: Not given.

Type locality: Not given.

Remarks: A form of Nerita fluviatilis (Figs. 37, 38).

Nerita quadricolor

Gmelin, J. F. 1791. Systema naturae, 13th edition,
p. 3684.

Description: N. testa subviolacea apice flavicante intus
alba striis elevatis nigris: labio utroque dentato,
exteriore intus sulcato; interiore rugoso.

Type locality: "Mari Rubro" (Red Sea).

Remarks: Species OK, Indian Ocean, resembles N. spengleri, see Mienis.

Nerita quoyi

LeGuillou, M. 1841. Description de quatorze nerites
nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 344.

Description: Testa globosa, solida, nigrata, albido-
maculata et unifasciata; anfractu ultimo superne
depresso; spira convexo-obtusa, apice erosos; apertura
rotundato-straminea, intus semi-lunari; labio plano
granuloso, margine tridentato; labro crasso, integro,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. guoyi, cont.

intus crenato, superne et inferne dentato.

Type locality: Unknown.

Nerita radiata

Recluz, M. C. A. 1841. Description de quelque nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 149.

Description: Testa globoso-acuta, sub epidermide lutescente albida, nigro late radiata, transversim sulcata; spira brevi, exsertiuscula, lutea, acuta; aperture alba, in fundo luteo; labio plano, concavo, laevi, margine dentibus 3 inequalibus; labro acuto, integro, intus calloso, obsolete sulcato, superne unidentato.

Type locality: "Je l'ai achetee avec des nerites noires et tres-noire (Ner. atrata et nigerrima)."

Nerita radula

Linnaeus, C. 1758. Systema naturae, p. 777.

Description: N. testa sulcata tuberculis aequalibus, labiis edentulis.

Type locality: Unknown.

Remarks: Now Neritopsis radula.

Nerita ramosa

Meuschen, F. C. 1787. Museum Geversianum, p. 170.

Remarks: Not acceptable.

Nerita rangiana

Recluz, M. C. A. 1841. Description de quelques especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 339.

Description: Testa minima, ovato-subhemisphaerica, laeviuscula, viridi, lineis seu punctis albis, fuscis, violaceisve variegata seu fasciata; anfractu infimo angustato, superne planulato, radiatim striato; spira brevissima; apertura subrotunda, anterius dilatata; labio convexo, albido, margine crenis quinis notato; labro superne anguloso sinuatoque.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. rangiana, cont.

Type locality: Madagascar.

Remarks: Now Smaragdina or Smargdella rangiana, see Japanese references.

Nerita rarispina

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 311.

Description: Testa minuta, globoso-ovata, rufo-nigra interdum maculis oblongis albidis notata; ultimo aufractu supra medium spinis nigris, raris 1-2-3 armato, spira exserta, apice erosa, peritremate rotundato, lutescente; columella plana, declivi, semilunari; margine inmedio subarcuata unidentata.

Type locality: "Hab. La Sicile, d'ou elle a ete rapportee par feu M. Caron, naturaliste-voyageur."

Nerita recluziana

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 345.

Description: Testa ovata, longitudinaliter rugosiuscula, fusco-nigricante, punctis irregularibus rufis undique ocellata; anfractibus ternis seu quaternis infimo superne angulosiusculo et spinifero; spira erosa, convexa; apertura dilatata, intus semi-lunari, labio luteo, plano, margine crenulato.

Type locality: Tahiti.

Remarks: "Neritine."

Nerita reticulata

Meuschen, R. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita reticulata

Meuschen, R. C. 1787. Museum Geversianum, p. 274.

Remarks: Not acceptable.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita reticulata

Karsten, D. L. G. 1789. Museum Leskeanum, Vol. 1, p. 296.

Description: N. ret. testa ovata, transversim profunde sulcata, longitudinaliter striata, striis exaratis tenuissimis et quam maxime adproximatis, unde facies retibus minutissimis composita, quin imo cancellata fere; apertura obovata denticulata; labrum marginé exteriori, subplicatum, interiori denticulatum dentibus obsolete compressis 13 labium rugosum, dentibus 2 marginalibus vix conspicuis rugescentibus.

Type locality: Not known.

Remarks: See ICZN Opinions.

Nerita reticulata

Quoy, J. R. C., and J. P. Gaimard. 1834. Voyage de decouvertes de l'Astrolabe..., Zoology, Vol. 3, p. 194.

Description: Nerita, testa ovato-convexa, levi, lutea, fusco aut nigro reticulata; spira obtusa, rotunda; apertura alba.

Type locality: "Habite les eaux douces du port Dorey, ja la Nouvelle-Guinee."

Nerita ringens

Reeve, L. A. 1855. Conchologia iconica: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 17, species 75.

Description: Ner. testa globosa, parva, solida, spira leviter exserta, anfractibus rotundatis, spiraliter regulariter costatis, labro et columella fortissime dentatis, area columellari tumida et corrugata; intense plumbea.

Type locality: Not given.

Nerita roissyana

Recluz, M. C. A. 1841. Description de quelques especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 338.

Description: Testa ovato-conoidea; subepidermide nigrescente, lineolis lutescentibus tenuissimis, angulato flexuosis, creberrimis, irregularibus notata; anfractibus quaternis vix convexis; apira exsertiuscula, conica,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. roissiana, cont.

obtustata; peristome ovato-acuto, intus flavo; columella angusta, subconvexa, in medio tenue arcuata et denticulata.

Type locality: New Guinea.

Nerita rossmassleriana

Recluz, M. C. A. 1845. Description de quelques nouvelles nerites fluviatiles, du cabinet de H. Cuming, Esq. Proc. Zool. Soc. London 13, p. 119.

Description: *Ner. testa subhemisphaerica, convexa, luteofulva, nigro irregulariter reticulata; anfractu unico laevissime perlongum striato; peritreme ovato-rotundato, basi obtuse anguloso; apertura crocea; columella planisima, declivi, margine acuto, in medio subarcuato et obsoletissime 3-4 denticulato.*

Type locality: Not given.

Nerita rubella

Muller, O. F. 1774. Vermium terrestrium et fluviatilium. Vol. 2, p. 195.

Description: *Nerita testa dilatata, convexa, nigricante, labio interiore crenulato.*

Type locality: "In Indiae fluviis."

Nerita rubra

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita rudis

Wood, W. 1828. Index testaceologicus, Supplement, p. 26.

Description: Not given.

Type locality: Africa.

Nerita rufa

*Born, I. von. 1778. Index Mus. Caes. Vind, p. 413. [not seen]

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita rugata

Recluz, M. C. A. 1842. Description de neuf especes de nerites nouvelles, suive d'observations sur les Nerita cornea et dubia. Rev. Zool. Soc. Cuvier. 1842, p. 75.

Description: Testa semiglolosa, ventricosa, superne vix depressa, rugoso-granulosa, nigra, inter oculum et luminem maculis subtriangularibus fuscis obsita; spira nullan apice rotundato-obtuso; apertura extus ovato-rotundata, albido-caerulescente; labio plano, declivi, margine inmedio tenuiter arcuato et obsolete rugoso.

Type locality: Philippines.

Remarks: Described in subgenus Clithon.

Nerita ruginosa

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 310.

Description: Testa semiglobosa, rugis angulato flexuosis, crebris arata; olivaceo-fusca, maculis triangularibus interdum picta; anfractibus ad carinam spinis armatis, ultimo basi costula transversa notato; apertura dilatata inferne oblique canaliculata et angulata; labio plano tenue arcuato et crenulato.

Type locality: Les iles sandwich, (M. Janelle).

Nerita rugosa

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3673.

Description: N. testa rugosa intus glaberrima dilute testacea: umbilici ambitu albo.

Type locality: "Habitat ad insulas Americae mediae obversas,"

Nerita rumphii

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 147.

Description: Testa ovato-subglobosa, transversim tenue sulcata, longitudinaliter tenue striata, posterius

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. rumphii, cont.

obsolete cancellata; maculis, fasciis seu flammulis picta; labio plano, basi vix rugoso et denticulato; labro intus tenuissime striato.

Type locality: "Hab. Amboine (Rumphius). La Nouvelle-Hollande, au port du roi Georges (Q. et Q.) l'ile Maria et les iles St-Pierre et St-Francois."

Remarks: Zoological Record 1877, Moll. p. 45 says rumphii synonym of N. polita (Pagenstecher); Vermeij says rumphii now doreyana, see Mienis.

Nerita samoensis

Dunker, G. 1871. Mollusca nova Musei Godeffroy Hamburgen-sis. Malakozool. Blatt. 18, p. 167.

Description: Testa parva, ovata, subglobosa, alba, fusco cinereoque maculata et variegata, vel unicolor atra, vel punctulis albis hinc illic adspersa, anfractibus transversim costatis et striatis instructa; costae striae que incrementi undulatae; spira exsertiuscula, apex saepius erosus; apertura latiuscula, interne sulcata; septum perparum cavatum et corrugatum in margine obsolete denticulatum, macula magna fusca notatum; operculum rugoso-granosum.

Type locality: Samoa.

Nerita sandalina

Recluz, M. C. A. 1842. Description de neuf especes de nerites nouvelles, suivie d'observations sur les Nerita cornea et dubia. Rev. Zool. Soc. Cuvier. 1842, p. 76.

Description: Testa semiglobosa, interdum subovata, subsutura compresso-strangulata, fusco-nigra, tenuiter striata; apice convexo-obtuso; apertura albido-lutescente; labio piano, margine vix arcuato et obsolete dentaticulato; labro patulo, superne depresso, linea nigra marginato.

Type locality: "Les iles Viti, a Sandal-bay, Lebouka, et les iles de la Societe a O-Tahiti."

Remarks: Described in the subgenus Clithon.

Nerita sanguinolenta

Menke, K. T. 1829. Berzeichniss de anfehnlichen

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. sanguinolenta, cont.

Conchylien-Sammlung des Freiherrn von der Malsburg, p. 15.

Description: Testa lateribus compressa, albo rubroque fasciata, sulcis inaequalibus etc.

Type locality: Not given.

Remarks: Red Sea species, similar to N. albicilla but smoother.

Nerita saturata

*Hutton, . 1884. Proc. Linn. Soc. N.S.W. 9(2), p. 354.
[not seen]

Remarks: N. saturata, new name for atrata (Lamarck, Reeve nec Chemnitz); in New Zealand atromentosa complex (Rehder).

Nerita savieana

Recluz, M. C. A. 1855. In L. A. Reeve. Conchologia iconica: or illustrations of the shells of molluscous animals 9 (Nerita), plate 8, species 37.

Description: Ner. testa globosa, crassissima, spira exserta, anfractibus convexis, latiusculis, spiraliter elevato-striatis et liratis, labro intus valde incrassato, dentato et crenulato, columella tumidiuscula, granato-corrugata, et dentata; caeruleo-nigra, interdum fluvo obscure variegata, intus flavicante.

Type locality: New Ireland.

Remarks: Recluz, MS Mus. Cuming.

Nerita sayana

Recluz, M. C. A. 1844. [untitled]. Proc. Zool. Soc. London 11, p. 199.

Description: Ner. testa ventricoso-ovata, longitudinaliter substriata, tenuiuscula, nitidula, nigrata, maculis fuscis seu lutescentibus, diversiformibus, variegata aut fusca et nigrescente nebulata, anfractibus duobus; primo partim derosos, infimo superne depresso; aperturea subrotunda, intus alba; labio calloso, plano, declivi, margine recto, in medio tenuissime crenato.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. sayana, cont.

Type locality: "Island of Guimaras, Philippines; in small streams. (H. Cuming)."

Nerita scabrella

Philippi, A. R. 1848. *Testaceorum novorum centuria*. Z. Malakozool. 5, p. 14.

Description: N. testa ovato-globosa, apice obtusissima, albida, nigro varia, transversim costata; costis ultra 30, irregulariter subgranosis; apertura extus suboviculari; labio concavo extus sulcato, centro granulato; margine columellari denticulis quatuor minimis munito; labro intus denticulis 14.

Type locality: Not given.

Nerita scabricosta

Lamarck, M. C. 1822. *Histoire naturelle des animaux sans vertebres*, vol. 6, part 2, p. 194.

Description: N. testa solida, transversim costata: costis elevatis, angustis, dorso asperulatis, nigris; interstiis albis; spira brevissima; apertura ringente, utriusque valde dentata; labio supra rugoso.

Type locality: "Mon cabinet."

Remarks: Original thought to be Indo-Pacific, now equated to Gulf of California species or subspecies as N. ornata or N. s. ornata (see Keene).

Nerita scalpta

Reeve, L. A. 1855. *Conchologia iconica*: or, illustrations of the shells of molluscous animals 9 (Nerita), plate 7, species 31.

Description: Ner. testa globosa, solidiuscula, spira sub-exserta, anfractibus laevibus, lineis impressis subdistantibus spiraliter incisis, labro intus lirato, columella granulata, margine tenue dentata; nigra, lineis impressis albidis, area columellari pallidissime lutescente.

Type locality: Not known.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita semiconica

Wood, W. 1828. Index testaceologicus, Supplement, p. 26.

Description: None given

Type locality: Not given.

Nerita semirugosa

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 102.

Description: N. testa maxima, semi-globosa supra depres-siuscula, crassa, luteo-rufescente, transversim sulcata, nigro trifasciata et undata; spria brevissima, prominula, acuta; apertura rotundata, dilatata, alba; labio lato, convexo, semilaevi, semirugoso, margine inaequaliter tridentato, lutescente; labro, intus striato, superne subbidentato.

Type locality: l'Ile-de-France selon Favanne, Timor et la Nouvelle-Hollande (Mus. Paris).

Nerita senegalensis

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3686.

Description: N. testa ovata obtusa solida nigerrima intus alba sulcata: labio utroque dentato.

Type locality: Senegalia.

Remarks: Valid name for west Africa (Senegal to Angola) species, see Nickles (1950).

Nerita serpentina

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 16.

Remarks: Nomen nudum, no valid standing.

Nerita serratilinea

*Mergerla, . In C. T. Menke. 1829. Syst. meth. Moll., p. 27. [not seen]

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita serrulata

Recluz, M. C. A. 1842. Description de neuf especes de nerites nouvelles, suivie d'observations sur les Nerita cornea et dubia. Rev. Zool. Soc. Cuvier. 1842, p. 76.

Description: Testa semiglobosa, ventricosa, longitudinaliter tenue striata, fusco-nigra, maculis luteo-fuscis diversimode picta; anfractibus superne depressis; spira conico-depressa; apertura albido-fuscescente, patula; labio plano, margine recto, serrulato.

Type locality: Sumatra.

Remarks: Described in subgenus Neritina.

Nerita signata

Lamarck, M. C. 1822. Histoire naturelle des animaux sans vertebres, vol. 6, part 2, p. 195.

Description: N. testa parvula, scabriuscula, transversim costata et striata, albo-lutescente, maculis spadiceis variegata; costis squamoso-scabris; spira incumbente, sublaterali; aperturae labiis minute dentatis; labio macula sanguinea notato.

Type locality: Nouvelle-Hollande (ex D. Macleay).

Remarks: Species now N. reticulata.

Nerita siquijorensis

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 198.

Description: Ner. testa ovato-transversa, prostice angustata, dorso-convexa, superne planulata, solida, longitudinaliter tenuiter et crebre striata, spadiceo-reticulata, interstitiis maculis oblongis, antice actuis, albidis seu lutescentibus picta; anfractibus 1 1/2; apice retuso; apertura extus ovata, intus lutescente; labio piano, margine et in medio vix arcuato et tenuiter crenato; labro subcontinuo, lateralibus rectiusculo.

Type locality: "Isle of Siquijor, in a small stream." H. Cuming [Philippines].

Remarks: Described in Sect. A. Gen. Neritina.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita smithii

Wood, W. 1828. Index testaceologicus, Supplement, p. 25.

Description: Not given.

Type locality: Not given.

Nerita sobrina

Recluz, M. C. A. 1845. Description de quelques nouvelles nerites fluviatiles, du cabinet de H. Cuming, Esq. Proc. Zool. Soc. London 13, p. 119.

Description: Ner. testa globoso-ovata, lutescente, lineis nigris distantibus latiusculis undatis vel angulato-flexuosis fulgurantibusque picta; anfractibus tribus, ad suturam nigro-marginatis; apira semishpaerica; columella convesiuscula, superne plus minusve callosa, inferne compresso-subcanaliculata, margine recta, vix denticualta; labro semi-elliptico, intus incrassato.

Type locality: Cayenne?

Remarks: This is most likely a Neritina.

Nerita solium

Recluz, M. C. A. 1845. Description de quelques nouvelles nerites fluviatiles, due cabinet de H. Cuming, Esq. Proc. Zool. Soc. London 13, p. 120.

Description: Ner. testa ovato-ventricosa, olivacea, postice maculis rugis triangularibus transversisque obsolete picta; anfractibus 1 1/2; infimo supra medium vix angulos, superne depresso; apice punto excavato notato; apertura alba, margine pallide lutescente sive aurantia; columella plano-compressa, superne callosa, margine rugosa et in medio excavata.

Type locality: Bencoonet, isle of Sumatra; in a muddy stream.

Remarks: Described in genus Clithon.

Nerita souleyetana

Recluz, M. C. A. 1842. Description de plusieurs especes de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 182.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. souleyetana, cont.

Description: *T. ventricosa-globosa*, *tenuiter striata*, *nitida*, *nigra*, *lineis longitudinalibus stemineis saepius undulatis ornata*; *anfractibus tribus*; *ultimo superne spinis maculatis instructo*, *spira prominula*, *apice decorticata*; *apertura rotundata*, *basi vix dilatata*, *flavicante*; *labio angusto*, *plano*, *medio concavo*, *superne calloso*, *antice subrecto*, *unidentato et obsolete ruguloso*.

Type locality: Not given.

Nerita sowerbiana

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 174.

Description: *N. testa subglobosa*, *tenue striata nigro et albido alternatim fasciata*; *fasciis nigris albo-articulatis*, *albidis maculis nigro-marginatis antice sagittatis ornata*; *anfractibus tribus*, *convexis*; *spira vix prominula*, *rotundata*; *apertura obliqua cinerascente*; *labio subconvexo*, *lutescente*, *antice denticulato arcuatoque*.

Type locality: Sinait, province of North Ylocos, island of Lucon: on the stony bed of a small river.

Nerita spadicea

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3672.

Description: *N. testa subglobosa solida spadicea subtus alba*, *apice caerulecente*, *faucis fasciis lateritiis*, *dorsi alba*.

Type locality: Habitat ad littora insulae S. Mauritii.

Nerita speciosa

*Sowerby, J. DeC. 1840. Trans. Geol. Soc. London, 2, expl. pl. 57. [not seen]

Nerita spengleriana

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 201.

Description: *Ner. testa orbiculato-conica*, *transversim laevissime sulcata*; *sulcis basi obsoletis*, *albida*,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. spengleriana, cont.

nigro-zonata; spira conico-acuta, lutescente, nigro-punctata, sulcis profundioribus; labio albo, laevigato, margine subbidentato; labro intus incrassato, laevissimo. Operculum pallide fuscum, tenuissime granulosum, subtus pallide rufum. Dente apicali obsoleto, infimo arcuato, transversim substriato, postice superficie dilatata et truncata.

Type locality: "Iloilo, isle of Panay; under stones at low water." H. Cuming.

Remarks: Valid species.

Nerita sphaerica

Muller, O. F. 1774. Vermium terrestrium et fluviatilium, Vol. 2, p. 170.

Description: Nerita testa globosa, cornea, vertice obtuso.

Type locality: Not given.

Nerita spinifera

Recluz, M. C. A. 1842. Description de plusieurs especes de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 183.

Description: T. globoso-acuta, tenuiter striata, griseo-nigrescente, punctulis albido triangularibus obsolete aspersa; anfractibus quaternis: ultimo ventricoso, ad suturam depresso, infra obtuse-angulato et spinis brevibus armato; spira conica, laevigata, caeruleo-nigra, acuta; apertura luteola; labio angusto, inaequaliter compresso-plano, denticulato et medio arcuato.

Type locality: L'ile de Guaham (Guam?).

Remarks: Described in subgenus Clithon.

Nerita spinosa

Wood, W. 1828. Index testaceologicus, Supplement, p. 25.

Description: Not given.

Type locality: Not given.

TABLE I. (Continued) CATALOG OF RECENT NERITA SPECIESNerita spinula

Bolten, J. F. 1798. Museum Boletnianum, Part 2, p. 16.

Remarks: Nomen nudum, no valid standing.

Nerita squamaepicta

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 169.

Description: N. testa ovato-subhemishpaerica, supra medium vix depressa, striis longitudinalibus remotis superne et antice subarcuatis impressa, sub epidermide olivaceo maculis squamiformibus antice nigro-marginatis undique pircata; apice rotundato-obtuso; apertura extus ovata, intus albido-caerulescente; labio plano, lutescente, medio albo, antice tenue arcuatim denticulato.

Type locality: Agoo, province of Pangasinan, isle of Lucon; in a rapid river.

Remarks: Now a species of Neritina; specimen collected at Guam by Vermeij sent to Meiniis.

Nerita squamulata

LeGuillou, M. 1841. Description de quatorze nerites nouvelles. Rev. Zool. Soc. Cuvier. 1841, p. 344.

Description: Testa globoso-obtusa, crassa, albido transversim costulatata longitudinaliter oblique striolata et squamosiuscula, spira retusa; apertura lutescente, rotundato-dilatata, intus semi-lunari; columella plano-concava, rugosa, granulata, et quadridentata; labor tenue crenato intus sulcato.

Type locality: Not given.

Nerita squarrosa

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 174.

Description: N. testa ventricoso-globosa, interdum sub-ovata, superne depressa et subangulata, epidermide squarrosa, maculis rufescentibus angulatis confertis nigro deu olivaceo marginatis picta, apice rotundata, apertura extus subquadrata, intus albido-caerulescente; labio supra

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. squarrosa, cont.

lutescente, piano, antice ruguloso, medio arcuato.

Type locality: Jimamailan, isle of Negros: on stone in a mountain stream.

Nerita stella

Dillwyn, L. W. 1817. A descriptive catalogue of recent shell, arranged according to the Linnaean method; ..., p. 1004.

Description: Shell sub-globular, transversely ribbed, with the summit obtuse and radiated; lips denticulated.

Type locality: East Indian seas.

Remarks: See Cernohorsky for notes, stella Gmelin right name for squamulata LeGuillou, see Gmelin from Chemnitz (not in Gmelin, 1791).

Nerita stellatus

Martyn, T. 1789. Universal conchologist, vol. 3, fig. 110.

Remarks: Invalid, rejected name.

Nerita stercusmuscarum

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3673.

Description: N. testa laevi nivea maculis punctisque subtilissimis rufis varia; umbilico gibbo bifido.

Type locality: "In mari mediterraneo et americano."

Nerita striata

*Burrow, E. J. 1815. Elem. conch., p. 172. [not seen]

Remarks: N. striata Burrow good name for mangrove species in N. undata complex.

Nerita strictaNerita strigilata

Quoy, J. R. C., and J. P. Gaimard. 1834. Voyage de decouvertes de l'Astrolabe, Vol. 3, p. 197.

Description: Nerita, testa ventricosa-oblonga, laevi,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. strigilata, cont.

nitida, strigis longitudinalibus alterne nigris et albis picta; spira exertiuscula, acuta; apertura alba; labio denticulato (Lamk.).

Type locality: Not given.

Remarks: "Cette Nerite, qui est encore fort recherchée, provient de Bourou, dans les Moluques. Elle se trouve aussi à Guam. L'exemplaire de ce dernier lieu est plus délicatement strié." p. 198.

Nerita striolata

Recluz, M. C. A. 1841. Description de quelques espèces nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 337.

Description: Testa globoso-obtusa, nitida, pallide olivacea, striolis undulatis transversis, regularibus, creberrimisque, sublente insculpta; anfractibus quaternis, superne lineola nigra interrupta submarginatis; spira vix exserta, apice erosa; labio obliquo, convexo, calloso, margine recto ac tenue ruguloso.

Type locality: Les Antilles?

Nerita subalata

Souleyet, M. 1842. Description de cinq nouvelles espèces de nerites fluviales provenant de voyage de la Bonite. Rev. Zool. Soc. Cuvier. 1842, p. 269.

Description: Testa ovata, transversa, tenui, convexa, fusco-nigricante, maculis parvis albidis suboculo et lumine tessellata; apertura postice oblique truncata, fusco-nigricante, labio antice arcuatim denticulato; denticulis retro sulco marginatis; labro postice superne et inferne subdilatato.

Type locality: Luzon, Philippine Islands.

Nerita subauriculata

Recluz, M. C. A. 1842. Descriptions of new species of nerites, collected by Mr. Cuming in the Philippine Islands. Proc. Zool. Soc. London 10, p. 171.

Description: N. testa ovato-hemisphaerica, convexa, fusco-violascente, punctis albido-lutescentibus dense aspersa; apice brevissimo; apertura extus ovata, postice truncata,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. subauriculata, cont.

continuo, plumbeo-nigrescente, dilatata; columella convexo depressa, sulco obsoleto circumcincta, anice vix arcuata, denticulata; labro superne in auricula brevissima pro-longato.

Type locality: Jaimamailan, isle of Negros: on the Nepa Palm trunks in brackish water.

Nerita subpunctata

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 199.

Description: Ner. testa semiglobosa, olivaceo-fusca, tenuiter rugata; rugis punctis nigrescentibus seriatis notatis, subepidermide alba, nigro reticulata; anfractu unico, supra medium angulosus; apice perforato; apertura extus subrotunda, intus albo-cinerascente; labio semi-lunari, plano, superne calloso, margine in medio tenuiter arcuato et crenulato; dente cardinali majore truncata.

Type locality: Sinait, province of North Ilocos, isle of Luzon; in a small mountain-stream. H. Cuming.

Remarks: Described in subgenus Clithon.

Nerita succincta

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

Nerita succinea

Recluz, M. C. A. 1841. Description de quelques especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 343.

Description: Testa minima, tenui, pellucida, ovato-semiglobosa, luteo-succinea, nitida; spira laterali, punctiformi; columella plana, margine recto, acuto.

Type locality: Madagascar, La Guadaloupe (Mus. Paris).

Nerita sulcata

*Born, I. von. 1778. Index Musei Caesarei vondobonensis, ..., p. 416. [not seen]

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita sulcosa

*Grateloup, . 1814. Conch. subop., p. 198. [not seen]

Nerita tenebrosa

Recluz, M. C. A. 1842. Description de plusieurs especes de nerites nouvelles vivantes. Rev. Zool. Soc. Cuvier. 1842, p. 180.

Description: Testa semiglobosa, crassa, nigerrima, punctulis albis adspersa; costis transversis 25-26 subaequilibus sculpta; spira brevissima, apice prominula et acuta; apertura estus rotundata, patula, nigro marginata; intus albida, angustata; labio plano, angusto, bidentato et in medio scrobicula notato; labio intus incrassato, breviter crenulato.

Type locality: "l'ile Solo, au nord de Borneo (M. Farges)."

Nerita tessellata

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3685.

Description: N. testa obtusa transversim striata: striis maculis subquadratis albis nigrisque, labio utroque denticulato.

Type locality: "Habitat ad insulas inter Africam et medium Americam intersitas,..."

Remarks: Valid West Indian species.

Nerita textilis

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3683.

Description: N. testa alba lineis angulatis nigris varia: costis sulcisque sedecim crenatis: labio exteriore exterius crenato intus dentato, interiore superne rugoso inferne tuberculato; palato dentato.

Type locality: Not given.

Remarks: Valid Indian Ocean species.

Nerita tigrina

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. tigrina, cont.

Remarks: Not acceptable.

Nerita tomolini

Turton, W. H. 1933. The marine shells of Port Alfred,
South Africa, errata page.

Remarks: Nomen novum for N. dubia Turton, 1932, a name pre-
occupied by N. dubia Recluz, 1842.

Nerita touranesnsis

Souleyet, M. 1842. Description de cinq nouvelles especes
de nerites fluviales provenant du voyage de la Bonite.
Rev. Zool. Soc. Cuvier. 1842, p. 269.

Description: Testa subhemisphaerica, postice angustata,
convexa, subepidermide olivacea albida, transversum 3-4
zonata: zonis lineis nigris intextis ornatis; spira
brevissima oblique incurba ac adpressa; apertura orbicu-
lari, continua albida; labio planiusculo, antice arcuato
et denticulato: denticulis pone fossula notatis.

Type locality: Tourane (Cochinchine).

Nerita trabalis

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita transversalis

*Ziegler, . In C. Pfeiffer. 1828. Naturg. deutsch.
land-moll., vol. 3, p. 48. [not seen]

Nerita transversaria

*M'Clelland, J., 1841. J. Nat. Hist., vol. 6, p. 243. [not
seen]

Nerita triangularis

Meuschen, F. C. 1787. Museum Geversianum, p. 174.

Remarks: Not acceptable.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita tricolor

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3686.

Description: *N. testa rubro nigro albidoque varia: anfractu primo striis quindecim elevatis, labio utroque dentato.*

Type locality: "Habitat ad Africae littus occidentale,..."

Remarks: See Cernohorsky.

Nerita trifasciata

LeGuillou, M. 1841. *Description de quatorze nerites nouvelles*. *Rev. Zool. Soc. Cuvier*. 1841, p. 343.

Description: *Testa ovata subglobata transversim sulcata sulcis 30-33 angustis, profundis; albido-viridescente, nigro trifasciata et sub flammulata; spira brevi, laterali, su acuta; labio plano, rugoso et granuloso, margine tridentato, dentibus medianis parvulis; labro intus incrassato, breve, sulcato, unidentato.*

Type locality: Triton-Bay, Nouvelle-Guinee.

Remarks: "Cette espece est voisine de la *N. chrysostoma* de M. Recluz..."

Type locality: Not given.

Nerita tritonensis

LeGuillou, M. 1841. *Description de quatorze nerites nouvelles*. *Rev. Zool. Soc. Cuvier*. 1841, p. 345.

Description: *Testa ovata, viridescente, lineis longitudinaliter flavescentibus arcuatissimis radiata; spira omnino erosa, planissima; apertura obliqua, labio angustiusculo, piano, margine subarcuato unidentato; labaro rotundato, superne estesque depresso.*

Type locality: Triton-Bay, Nouvelle-Guinee.

Nerita trochus

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 176.

Description: *Nerita testa trochiformi, carinata, apertura coarctata.*

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. trochus, cont.

Type locality: "In Saxonia Weimariensi."

Nerita tuberculata

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*, Vol. 2, p. 191.

Description: *Nerita testa subulata, cinerea, transversim striata; anfractibus nodulosis, stirisque sanguineis.*

Type locality: "In littore Coromandel cum B. scabro frequens."

Nerita turrita

Gmelin, J. F. 1791. *Systema naturae*, 13th edition, p. 3686.

Description: *N. testa alternatim albo nigroque fasciata intus alba: vertice valde clata, labro acuta.*

Type locality: In insularum antillarum.

Remarks: Valid species as Neritina turrita, a mangrove species.

Nerita turtoni

Recluz, M. C. A. 1843. [untitled]. Proc. Zool. Soc. London 11, p. 71.

Description: *Ner. testa ventricoso-ovata, luteo-rufa seu rubicunda, flammis nigrescentibus undatis, remotisque, sive angulato-flexuosis picta; anfractibus 4-5, convexis, supremis derosis: infimo superne horizontaliter depresso; labio subconvexo, fusco-rubente, margine in medio tenuiter crenato, basi leviter emarginato; labro intus calloso-alba, ad marginem fusco rubente.*

Type locality: Not given.

Nerita umlaasiana

*Krauss, C. F. F. 1848. *Sudafrik. Moll.*, p. 89. [not seen]

Remarks: Valid South African and Madagascar species.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita undata

Linnaeus, C. 1758. *Systema naturae*, p. 779.

Description: *N. testa sulkata sulcis triginta, labiis dentatis: interiore rugoso tuberculato.*

Type locality: "Habitat in O. Asiae ad Amboinam."

Remarks: Valid species.

Nerita undulata

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3678.

Description: *N. testa tenui laevi undulata: vertice obtuso, labio exteriori substriato edentulo, interiori subdenticulato.*

Type locality: Not given.

Nerita urceus

Muller, O. F. 1774. *Vermium terrestrium et fluviatilium*,
Vol. 2, p. 174.

Description: *Nerita testa subumbilicata, ventricosa, rugosa, fusca; sperture essusa alba.*

Type locality: "In insulis Indiae edulis. In Museo Moltkiano."

Nerita usnea

Bolten, J. F. 1798. *Museum Boltenianum*, Part. 2, p. 16.

Remarks: Nomen nudum, no valid standing.

Nerita valvata

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3675.

Description: *N. testa planiuscula: apertura circinata.*

Type locality: "Habitat in Oceano septenatrionali,"

Nerita varia

Meuschen, F. C. 1787. *Museum Geversianum*, p. 268.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. varia, cont.

Remarks: Not acceptable.

Nerita variegata

Bolten, J. G. 1798. Museum Boltenianum, Part 2, p. 18.

Remarks: Nomen nudum, no valid standing.

Nerita versicolor

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

Remarks: Not acceptable.

Nerita versicolor

Gmelin, J. F. 1791. Systema naturae, 13th edition,
p. 3684.

Description: N. testa maculis quadratis nigricantibus,
caerulecentibus, rubris albisque et fasciis rubro alboque
maculatis varia: labio interiore intus striato utrinque
dentato.

Type locality: "Habitat ad insulas Antillas,..."

Remarks: Valid West Indian species.

Nerita vestita

Souleyet, M. 1842. Description de cinq nouvelles especes
de nerites fluviales provenant du voyage de la Bonite.
Rev. Zool. Soc. Cuvier. 1842, p. 269.

Description: Testa ventricoso-globosa, ad suturam depressa,
nitida, tenuiter striata, subepidermide nigerrima trans-
versim obscure zonata; spira brevi, apice erosa; apertura
caerulecente; labio plano, albido, antice edentulo; labri
epidermide incolentibus productiori.

Type locality: Luzon, Philippine Islands.

Nerita vexillum

Reeve, L. A. 1855. Conchologia iconica: or, illustrations
of the shells of molluscous animals 9 (Nerita), plate 16,
species 65.

Description: Ner. testa ovata, spira subexserta,

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. vexillum, cont.

anfractibus rotundatis, juxta suturam impressis, spiraliter regulariter liratis, columella concavo-plana, acute denticulata; fasciis latis nigris et luteis cingulata, nigro luteo-punctato.

Type locality: Not known.

Nerita violacea

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita violacea

Gmelin, F. J. 1791. Systema naturae, 13th edition, p. 3636.

Description: N. testa ovata violacea albo punctata solida laevi: labio interiori inferne denticulato.

Type locality: Not given.

Nerita virens

Meuschen, F. C. 1787. Museum Geversianum, p. 270.

Remarks: Not acceptable.

Nerita virgata

Meuschen, F. C. 1787. Museum Geversianum, p. 274.

Remarks: Not acceptable.

Nerita virginaea

Linnaeus, C. 1758. Systema naturae, p. 778.

Description: N. testa laevi ovata, labio interiore gibbo denticulato.

Type locality: "Habitat in M. Mediterraneo."

Remarks: Now Neritina virginaea, see Russell.

Nerita viridicata

Meuschen, F. C. 1787. Museum Geversianum, p. 268.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. viridicata, cont.

Remarks: Not acceptable.

Nerita viridis

Linnaeus, C. 1758. *Systema naturae*, p. 778.

Description: N. testa laevi viridi, labio interiore medio crenulato.

Type locality: Minorcam, Jamaica.

Remarks: Now Smaragdia [sp?] viridis from Mediterranean.

Nerita viridis

Meuschen, F. C. 1787. *Museum Geversianum*, p. 272.

Remarks: Not acceptable.

Nerita vitellus

Linnaeus, C. 1758. *Systema naturae*, p. 776.

Description: N. testa umbilicata subglobosa, umbilico perforato aequali.

Type locality: "Habitat in O. Asiae."

Nerita vittata

Gmelin, J. F. 1791. *Systema naturae*, 13th edition,
p. 3674.

Description: N. testa subglobosa fusca: vitta duplii alba
medio reticulata utrinque denticulata.

Type locality: "Habitat ad Marochii littora."

Nerita vivipara

Muller, O. F. 1774. *Vermium terrestrium et fluviatile*,
Vol. 2, p. 182.

Description: Nerita testa ovato-ventricosa, virescente,
fasciis tribus lividis, obscuris.

Type locality: Not given.

Remarks: Now considered a species of Helix.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESNerita webbei

Recluz, M. C. A. 1849. Description de quelques nouvelles especes de coquilles. Rev. Zool., Ser. 2, 1, p. 70.

Description: Testa subglobosa seu ovato-conica, superne convexo-obtusissima, lavigata, lutescente interdum pallide fusca, lineolis nigris, rubrisve, longitudinalibus aut transversis varie picta; anfractibus 1 1/2; labio incrassato, gibboso, margine in medio irregulariter denticulato; labro acuto, tenui.

Type locality: Grand-Bassam, west coast of Africa.

Remarks: Described in subgenus Neritina.

Nerita winteri

Philippi, R. A. 1844. Abbildungen und beschreibungen neuer oder wenig gekannter conchylien,..., p. 86.

Description: N. testa ovata, albida, transversim costato-sulcata; spira elevata, acuta; anfractu ultimo superne depresso; labro intus tenuissime denticulato; labio plano slucis aliquot exarato; columella quadridentata, dentibus medianis latis.

Type locality: Mollucas.

Nerita yoldii

Recluz, M. C. A. 1841. Description de quelques nouvelles especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 151.

Description: Testa parva, semiglobosa, obtusata, nigra, maculis flammisque angulato flexuosis alios, saepissime erosio-impressis insculpta; apertura rotundata, dilute straminea; labio rugoso granosoque, basi in medio marginae dentibus binis parvulis instructo; labro intus sulcato, superne obsolete dentato.

Type locality: Coast of Brazil.

Remarks: Valid(?) in Korea and China, Vermeij reported on S.I. specimen.

Nerita zebrina

Recluz, M. C. A. 1841. Description de quelques especes de nerites vivantes. Rev. Zool. Soc. Cuvier. 1841, p. 341.

TABLE 1. (Continued) CATALOG OF RECENT NERITA SPECIESN. zebrina, cont.

Description: Testa parvula, semiglobosa, tenui, laevissime striata, luteo-virescente, lineolis longitudinalibus rufo-purpurascenscentibus angulato-flexuosis picta; anfractibus 2-3 convexis: ultimo superne compresso; spira vix exserta; apertura rotundata, alba; labio plano, edentulo.

Type locality: "Les environs de Montpellier,..."

Nerita zelandica

Recluz, M. C. A. 1845. Description de quelques nouvelles nerites fluviatiles, due cabinet de H. Cuming, Esq. Proc. Zool. Soc. London 13, p. 120.

Description: Ner. testa ovato-oblonga, ventricosa, tenuiuscula; anfractibus 3-4 supremis saepius derosis, infimo subsutura horizontaliterque compresso; nigra lineis ravidis longitudinalibus angulato-flexuosis creberrimis picta, interdum lutescente superne et inferne late fasciata; columella subcompressa, crocea, margine denticulata et in mediovix arcuata; labro tenui margine sordide rubente, intus lacteo ac parum incrassato.

Type locality: New Zealand: on stones in mountain-streams.

Nerita ziczac

Meuschen, F. C. 1787. Museum Geversianum, p. 272.

Remarks: Not acceptable.

Nerita ziczac

Bolten, J. F. 1798. Museum Boltenianum, Part 2, p. 16.

Remarks: Nomen nudum, no valid standing.

TABLE 2

LIST OF FOSSIL NERITA SPECIES [* = not seen]Nerita abeihensis

Whitfield, R. P. 1891. Observations on some Cretaceous fossils from the Beyrut district of Syria, in the collection of the American Museum of Natural History, with descriptions of new species. Bull. Am. Mus. Nat. Hist. 3:431. [Abeih Sandstone, near Abeih, Syria]

Nerita acherontis

*Brongniart, A. 1823. Mem. Terr. Vicentin., p. 60.

Nerita alpina

*Keipstein, A. v. 1845. Beitr. Geol. Ostl. Alpen (2,3), p. 200. [Triassic fossil, not Nerita]

Nerita alternata

*Schleuter, F. 1838. Kurzg. Syst. Verz. Conch. 1838, p. 15. [nomen nudum, Index Animalium 2]

Nerita ampilata

*Koninck, L. G. de. 1844. Desc. Anim. Foss. (II) 1844, p. 485. [Natica, Index Animalium 2]

Nerita angistomoidea

*Korobkov, I. A. 1961. New genus of Eocene molluscs from southern Ukraine. Paleont. Zh. 1961(4), p. 34. [Eocene, southern Ukraine]

Nerita angistoma

Deshayes, G. P. 1832. Encyclopedie methodique, Vol. 3, p. 622.

Nerita angolensis

Rennie, J. V. L. 1929. Cretaceous fossils from Angola (Lamellibranchia & Gastropoda). Ann. South African Mus. 28, p. 39. [Cretaceous Angola]

Nerita angulata

*Sowerby, J. De C. 1836. Trans. Geol. Soc. London 4, p. 347.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita anodonta

*Melville, J. C. 1898. Further investigations into the molluscan fauna of the Arabian Sea, Gulf of Oman, and Persian Gulf, with descriptions of forty species. Mem. Manchester Soc. 42(4), p. 24. [Karachi]

Nerita bavarica

*Schiffhautl, K. E. 1854. Beitrage zur naheren Kenntniss der Bayern'schen Voralpen. Neue Jahrb. Mineral., Geogn., Geol., Petrefak. p. 524. [?Mesozoic-Triassic]

Nerita beali

Loel, W., and W. H. Corey. 1932. The Vaqueros formation, Lower Miocene of California. I. Paleontology. Univ. California Pub. Geol. 22(3), p. 272. [Miocene California]

Nerita benecki

*Rollier, L. 1918. Fossiles nouveaux ou peu connus des terrains secondaires (Mesozoiques) du Jura et des contrees environnantes. Mem. Soc. Pol. Suisse Geneve 43, p. 32. [Jurassic]

Nerita bergadana

*Vidal, D. L. M. 1921. Contribution a la paleontologia del cretacico de Cataluna. Mem. Roy. Acad. Di. Barcelona 17, p. 15.

Nerita biangulata

Dickerson, R. E. 1915. Fauna of the Martinez Eocene of California. Univ. California Publ. Bull. Dept. Geol. 8, p. 141. [California Eocene]

Nerita bicoronata

Deshayes, G. P. 1864. Description des animaux sans vertebres decouverts dans le bassin de Paris...3, p. 13.

Nerita bidens

Whitfield, R. P. 1891. Observations on some Cretaceous fossils from the Beirut district of Syria, in the collection of the American Museum of Natural History, with descriptions of some new species. Bull. Am. Mus. Nat Hist. 3, p. 432. [Abeih, Syria]

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita bisinuata

*Buvignier, A. 1842. In Sauvage and Buvignier. Stat. Min. Geol. Ardennes p. 535.

Nerita boriesi

*Doncieux, L. 1908. Ann. Univ. Lyon 22, p. 237.

Nerita bourgeoisiana

d'Orbigny, A. 1850. Prodrome de Paleontologie...2, p. 192.

Nerita brimonti

Deshayes, G. P. 1864. Description des animaux sans vertebres decouverts dans le bassin de Paris 3, p. 14.

Nerita broungniartina

d'Orbigny, A. 1850. Prodrome de Paleontologie...2, p. 301.

Nerita bronni

*Cristofori, ., and . Jan. 1832. Cat. Mus. Sect 2, p. 4.

Nerita brouzetensis

*Cossmann, M. 1921. Rectification de nomenclature. Rev. Crit. Paleonzool. 25, p. 79.

Nerita bruni

*Cossmann, M. 1916. Le barremien superieur a facies urgonien de Brouzet-les-Alais Gard. Part II. Complement de l'etude paleontologique des gisements de Brouzet. Mollusques (Gastropodes et Pelecypodes). Mem. Soc. Geol. France Paleontol. 21(51), p. 24. [Urgonian France]

Nerita bruni

*Roman, F., and P. Mazeran. 1920. Monographie paleontologique de la faune Turonien du Bassin d'Uchaux et de ses dependances. Lyon Arch. Mus. Hist. Nat. 12, p. 38. [Cretaceous France]

Nerita burdigalensis

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 40. [early Miocene, France]

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita caffra

Wood, W. 1828. Index testaceologicus, Supplement, p. 25.

Nerita cancellata

*Morris, J., and J. Lycett. 1854. A monograph of the Mol-lusca from the Great Oolite, chiefly from Minchinghamton and the coast of Yorkshire 1, p. 56. [Jurassic]

Nerita cancellata

*Zieten, C. H. v. 1832. Petrif. Wurt. 6, p. 44. [IA2 noted Neritites]

Nerita capduri

*Cossmann, M. 1907. Le barremien superieur a facies urgonien de Brouzet-les-Alais (Gard). Description des Gastropodes et Pelecypodes. Mem. Soc. Geol. 37, p. 22.

Nerita carapezzai

*Checchia-Rispoli, G. 1912. Sopra alcunimolluschi eocenici della Sicilia. Palermo G. Sci. Nat. Econ. 29, p. 95. [Eocene Sicily]

Nerita carolina

*Stoliczka, F. 1868. The Gastropoda 2, p. 341. [Cretaceous]

Nerita caronis

*Brongniart, A. 1823. Mem. Terr. Vicentin., p. 60.

Nerita cassidiformis

Lycett, J. 1850. Tabular view of fossil shwells from the Middle Devonian of the Inferior Oolite in Gloucestershire. Ann. Mag. Nat. Hist., Ser. 2, 6, p. 416. [Jurassic]

Nerita cenomanensis

*Gueranger, E. 1853. Essai d'un repertoire paleontologique du department fe la Sarthe...p. 31. [Cretaceous]

Nerita cestophora

*Ryckholt, P. 1852. Melanges paleontologiques 1, p. 82.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita circumvallata

*Bayan, F. 1870. Sur les tertiaraires de la Venetie. Bull. Soc. Geol. France, Ser. 2, 27, p. 458. [?Eocene]

Nerita clathrata

Gmelin, J. F. 1791. Systema naturae, 13th edition, p. 3675. [Fossils in Campania Gallica]

Nerita clavatula

*Lycett, J. 1863. Supplementary monograph on the Mollusca from the Stonesfield Slate, Great Oolite, Forest Marble, and Cornbrash, p. 98.

Nerita coffati

*Cox, L. R. 1925. Cretaceous Gastropoda from Portuguese East Africa. Ann. Transvaal Mus. 11, p. 208.

Nerita colpostoma

*Schafhautl, K. E. 1863. Sud-Bayerns lethaea geognostica, p. 183.

Nerita compacta

*Forbes, E. 1856. Report on the fossil invertebrata from southern India, collected by Mr. Kaye and Mr. Cunliffe. Trans. Geol. Soc. London 7, p. 122.

Nerita consobrina

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 31. [Neritina from Index Animalium 2]

Nerita constricta

*Loriol, P. de, and E. Koby. 1890. Etudes sur les Mollusques des Couches Coralligenes inferieures du Jura Bernois. Part I. Abh. Schw. Pal. Ges. 16, p. 105.

Nerita corallina

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 7.

Nerita corex

*Vokes, H. E. 1939. Molluscan faunas of the Domengine and

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

N. corex, cont.

Arroyo Hondo formation of the California Eocene. Ann. N. Y. Acad. Sci. 38, p. 180. [California Eocene]

Nerita corneti

*Vincent, E. 1930. Etudes sur les mollusques montiens du Pondingue et du Tuffeau de Ciply. Mem. Mus. Hist. Nat. Belgique 46, p. 20.

Nerita costellata

*Muesnster, . 1844. In G. A. Goldfuss. Petref. German 3(8), p. 115. [Triassic]

Nerita costulata

Deshayes, G. P. 1838. In J. B. P. A. de Lamarck. Histoire naturelle des animaux sans vertebres, Second edition, p. 617.

Nerita costulata

*Roemer, F. A. 1841. Verst. Norddeutsch. Kreidegeb. 2, p. 82.

Nerita cowlitzensis

*Dickerson, R. E. 1915. Fauna of the Type Tejon: its relation to the Cowlitz Phase of the Tejon Group of Washington. Pub. California Acad., Ser. 4, 5, p. 58. [Eocene North America]

Nerita crassa

*Bellardi, L. 1852. Catalogue raisonne des fossiles nummulitiques du Comte de Nice. Mem. Soc. Geol. France, Ser. 2, 4, p. 212.

Nerita crassecallosa

*Blaschke, F. 1911. Zur Tithonfauna von Stramberg in Mahren. Wien Ann. Nat. Hist. Hofmus. 25, p. ? [Stramberg Mahren Jurassic]

Nerita crenata

Anton, H. E. 1839. Verzeichniss der Conchylien...., p. 30.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita crooki

Clark, B. L. 1938. Fauna from the Markley formation (Upper Eocene) on Pleasant Creek, California. Bull. Geol. Soc. America 49(5), p. 700. [Upper Eocene, Pleasant Creek, California]

Nerita cubanensis

*Pcelinzev, V. 1928. On some Jurassic fauna from the Bolshie Balkhony (Turkmenia). Bull. Com. Geol. Leningrad 46, p. 1179. [in Russian, Cretaceous South Russia]

Nerita cuneata

*Gabb, W. M. 1864. Description of the Cretaceous fossils. Palaeontology of California 1, p. 137.

Nerita custugensis

*Doncieux, L. 1908. Ann. Univ. Lyon 22, p. ?. [no pages given]

Nerita decorata

*Muenster, G. v. 1841. Beitr. zur Petref. 4, p. 98. [Triassic]

Nerita deformis

*Gabb, W. M. 1864. Description of the Cretaceous fossils. Palaeontology of California 1, p. 137,

Nerita denticulata

Stephenson, L. W. 1947. New Upper Cretaceous fossils from Mississippi and Texas. Part 1. Fossils from two deep wells in Missouri. Part 2. A new Venericardia from Uvalde County, Texas. Prof. Pap. U.S. Geol. Surv. 210E, p. 181.

Nerita deshayesea

*Buvignier, A. 1852. Statistique geologique, mineralogique et paleontologique du department de la Meuse, Atlas p. 30.

Nerita distorta

*Hornes, M. 1856. Die fossilen mollusken des Tertiaer-Beckens von Wien unter der Mitwirkung von Paul Partsch. Abhandlung. kaiser.-konigl. Geol. Reichsan. 3, p. 537.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita fajumensis

*Oppenheim, . 1906. Palaeontographica 30, p. 227.
[Tertiary Egypt]

Nerita favarottaensis

*Gemmellaro, G. G. 1866. Naticidae e Neritidae del terreno giurassico del mord di Sicilia. G. Sci. Nat. Econom. 2, p. 182.

Nerita fittoni

d'Orbigny, A. 1850. Prodrome de Paleontologie...2, p. 68.
[Neritina]

Nerita funata

*Dujardin, F. 1837. Mem. Soc. Geol. France 2, p. 417.

Nerita gaasensis

*Magne, A., and A.-M. Vergneau-Saubade. 1974. Les Neritiidae tertiaires de l'Aquitaine occidentale. Bull. Soc. Borda 356, p. 485. [Lower Oligocene Gaas-Lesbarritz, France]

Nerita gabrieliani

*Aslanyan, P. M. 1959. New species of Mollusca in Palaeogene deposits of south-western Armenia. Bull. Acad. Sci. Armenia SSR (Geol. Geog.) 12, p. 20. [Palaeogene, SW Armenia, in Russian]

Nerita galloprivinalis

*Matheron, P. 1843. Catal. meth. foss. B. du Rhone, p. 227. [ALSO Rep. Trav. Soc. Stat. Marseille 4, p. 299]

Nerita gea

d'Orbigny, A. 1852. Prodrome de paleontologie...2, p. 232.

Nerita gemmata

*Petho, J. 1906. Die Kraide-(Hypersenon-) Fauna des Perterwardeiner (Petervarader) Gebirges (Fruska Gora). Palaeontographica 52, p. 126.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita duboisi

*Rollier, . 1918. Mem. Soc. Pol. Suisse 43, p. 32.
[Jurassic]

Nerita duplicata

Wood, W. 1828. Index testaceologicus, Supplement, p. 25.

Nerita elegans

d'Orbigny, A. 1850. Prodomus de paleontologie...2, p. 346.
[Neritina]

Nerita equina

*Bezancon, A. 1870. Descriptions d'espèces nouvelles du bassin de Paris. J. Conchyliol. 18, p. 320.

Nerita etalloni

*Maire, V. 1926. Etudes géologiques et paleontologiques sur l'arrondissement de Gray. Les gastropodes du jurassique supérieur graylois. Bull. Soc. Grayloise 19, p. 96.

Nerita eucleista

*Oppenheim, . 1906. Palaeontographica 30, p. 288.
[Tertiary, Egypt]

Nerita euxina

*Clessin, S. 1886. Binnenmollusken aus Rumanien. Mala-kozool. Blat. 7, p. 49. [late Miocene, Bulgaria]

Nerita expansa

*Reuss, A. E. 1856. In M. Hornes. Die fossilen mollusken de Tertiaer-Beckens von Wien unter der Mitwirkung von Paul Partsch. Abhandlung. Kaiserlich-königlichen Geol. Reichsan. 3, p. 536.

Nerita exuviooides

*Trechmann, C. T. 1935. The geology and fossils of Carriacou, West Indies. Geol. Mag. 72, p. 551. [Oligocene Carriacou]

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita gensaciensis

*Petho, J. 1906. Die Kraide-(Hypersenon-) Fauna des Perterwardeiner (Petervarader) Gebriges (Fruska Gora). Palaeontographica 52, p. 117. [Upper Cretaceous]

Nerita gigantea

*Bellardi, L., and G. Michelotti. 1841. Mem. Roy. Acc. Sci. Torino 3, p. 164.

Nerita glaucinoides

*Morris, J. 1843. Cat. Brit. Foss., p. 154. [Natica]

Nerita glebosa

*Ryckholt, P. de. 1852. Melanges paleontologiques 1, p. 81.

Nerita giberti

Meinis, H. K. 1975. Note on recent and fossil Neritidae. 6. Nerita giberti, new name for Nerita morio Dujardin, 1837, non Nerita morio (Sowerby, 1832). Basteria 39, p. 1.

Nerita globosa

Sowerby, J. DeC. 1823. Min. Conch., vol. 5, p. 29.

Nerita globulus

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 301.

Nerita granulata

*Hoeninghaus, F. W. 1831. Jahrb. Min., p. 139. [nomen nudum]

Nerita granulosa

*Deshayes, G. P. 1832. Desc. Coq. Foss. Paris, 2, p. 159 [?p. 622]

Nerita granulosa

*Cristofori, ., and . Jan. 1832. Cat. Mus. (Sect. II) Conch. foss., p. 4.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita grateloupiana

*Ferussac, A. E. 1823. In G. P. Deshayes In A. E.
Ferussac. H. N. g. et. p. Moll., wrapper of liver. 20.
[n. et f.]

Nerita grossa

*Bronn, H. G. 1848. Index Pal., p. 805. [Neritites]

Nerita grovana

*Ferussac, A. E. 1823. In G. P. Deshayes In A. E.
Ferussac. H. N. g. et. p. Moll., wrapper of liver 20.
[n. et f.]

Nerita haliotis

*Sowerby, J. DeC. 1839. In R. I. Murchinson. Silurian system, p. 625. [not neritacean if Silurian]

Nerita hantkeni

*Szoto, E. 1953. Mollusques Eocenes de la hongrie. I. Les mollusques Eocenes de environs de Gant. Geol. Hung. 22, p. 16. [French and Russian summaries]

Nerita hantoniensis

Pilkington, W. 1804. Descriptions of some fossil shells found in Hampshire. Trans. Linnean Soc. London 7, p. 118.

Nerita helicina

*Brocchi, G. 1814. Conch. Subsp., p. 297.

Nerita hemisphaerica

*Roemer, F. A. 1836. Verst. Norddeutsch. oolith. 3, p. 156.

Nerita hieroglyphica

*Wattebled, G. 1886. Description de mollusques inedits de l'Annam. Recolte du capitaine Dorr aux environs de Hue. J. Conchyliol. 26, p. 68.

Nerita hilarula

*Nardo, G. D. 1847. Sin. mod. spec. Lag. Veneto, p. 97.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita hisingeri

*Bellardi, L., and G. Michelotti. 1841. Mem. Roy. Acc. Sci. Torino 3, p. 165.

Nerita hoffmanni

*Gemmellaro, G. G. 1866. Naticidae e Neritidae del terreno giuasico del nord di Sicilia. G. Sci. Nat. Econom. 2, p. 182.

Nerita ickei

*Martin, V. 1916. Die altnoicane Fauna des West-Progogebirges auf Java. A. Gastropoda. Leiden. Samml. Geol. Reichsmus., N.F., 2, p. 258.

Nerita incrassata

*Gemmellaro, G. G. 1866. Naticidae e Neritidae del terreno giuasico del nord dei Sicilia. G. Sci. Nat. Econom. 2, p. 181.

Nerita inflata

*Piette, E. 1855. Observations sur les etages inferieurs du terrain jurassique dans les departements des Ardennes et de l'Aisne. Bull. Soc. Geol. France, Ser. 2, 12, p. 1110.

Nerita inornata

*Piette, E. 1855. Observations sur les etages inferieurs du terrain jurassique dans les departements des Ardennes et de l'Aisne. Bull. Soc. Geol. France, Ser. 2, 12, p. 1110.

Nerita intermedia

*Grateloup, . 1828. Bull. H. N. Soc. Linn. Bordeaux, 2, p. 148.

Nerita intermedia

*Deshayes, G P. 1832. In C. Belanger, Voy. Indies Orient 6, p. 420.

Nerita involuta

*Lycett, J. 1863. Supplementary monograph on the Mollusca from the Stonesfield Slate, Great Oolite, Forest Marble, and Cornbrash, p. 20.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita ishidae

*Masuda, K. 1966. Molluscan fauna of the Higashi-Innai Formation of Noto Peninsula, Japan. II. Remarks on molluscan assemblage and description of species. Trans. Proc. Palaeontol. Soc. Japan 64, p. 335.

Nerita jaculatrix

*Schrantz, F. v. P. 1803. F. Boica 3(2), p. 290.

Nerita jaequinensis

*Addicott, W. O. 1970. Miocene gastropods and biostratigraphy of the Kern River area, California. Prof. Pap. U.S. Geol. Surv. 642, p. 45.

Nerita jayanica

*Olsson, A. A. 1944. Contribution to the paleontology of northern Peru, Part VII. The Cretaceous of the Paita Region. Bull. American Paleontol. 28(3), p. 17.

Nerita jurensis

*Muenster, G. v. 1836. Verz. Verst. Nat.-Samml. Baireuth, p. 50.

Nerita kaffraria

*Woods, H. 1906. The Cretaceous fauna of Pondoland. Ann. South African Mus. 4(7), p. 311.

Nerita kamigiriensis

*Itoigawa, J. 1957. On the Miocene Lucinidae from the Mizunami Group, Japan. Trans. Palaeontol. Soc. Japan 25, p. 140.

Nerita klipsteini

*Hornes, M. 1855. Über die Gastropoden und Acephalen der Hallstatter Schichten. Dekschr. Kaiserlichen Akad. Wissenschaft. 9, p. 41.

Nerita laevigata

*Sowerby, J. 1821. Min. Conch. 3, p. 31.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita laffoni

*Merian, P. 1846-48(1849). Ber. Nat. Ges. Basel 7, p. 31.

Nerita lamarmorae

*Gemmellaro, G. G. 1866. Naticidae e Neritidae del terreno giurassico del nord di Scilia. G. Sci. Nat. Econom. 2, p. 181.

Nerita laufonensis

*Thurman, J. 1862. In J. Thurman and A. Etallon. Letheia Bruntrutana ou etudes paleontologiques et stratigraphiques sur le Jura Bernois et en particulier les environs de Porrentruy, p. 121.

Nerita liasina

d'Orbigny, A. 1850. Prodrome de paleontologie...1, p. 214.
[Neritina]

Nerita libera

*Nardo, G. D. 1847. Lin. Mod. Spec. Lag. Veneto, p. 97.

Nerita littoralis

*Risso, A. 1826. H. N. Europe, 4, p. 151.

Nerita lochensis

*Brosamlen, R. 1909. Beitrag zur Kenntnis der Gastropoden des schwabischen Jura. Palaeontographica [Stuttgart] 56, p. 244.

Nerita maculata

*Doncieux, L. 1908. Ann. Univ. Lyon 22, p. 236.

Nerita mais

*Buvignier, A. 1843. Mem. Soc. Philom. Verdun 2, p. 241.

Nerita major

*Sandberger, F. 1863. Die Conchylien des Mainzer Tertiärbeckens, p. 158.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita mammaeformis

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 104.

Nerita mammillaris

*Thunberg, C. P. 1791. D. D. Mus. Ac. Upsal., App. 2,
p. 129. [Helix from Index Animalium 1]

Nerita marina

*Chiaje, S. d. 1830. Mem. An. s. Vert. Napoli, Tab. pro V
& VI, pl. 104.

Nerita marmorata

*Link, H. F. 1807. Bersch. Nat. Samml. Univ. Rostock 3,
p. 118.

Nerita martiniana

*Matheron, P. 1843. Catal-meth. foss. B. du Rhone, p. 228;
Rep. trav. Soc. stat. Marseille 6, p. 300.

Nerita martiniana tricostata

*Kecsheimetine-Kormendy, A. 1962. Neue Molluskenarten aus
dem Mittelmiozan von Varpalota. I. Gastropoden. Foldt.
Kozl. 92, p. 84. [Magyar with German summary; Varpalota,
Hungary, Miocene]

Nerita matonia

*Risso, A. 1826. H. N. Europe 4, p. 271.

Nerita meridionalis

*Philippi, R. A. 1836. Enum. Moll. Siciliae 1, p. 159.

Nerita michaillensis

*Pictet, F. J., and G. Campiche. 1862. Description des
fossiles du terrain cretace des environs de Sainte-Criox.
Materiaux pour la paleontologie Suisse, Ser. 3, p. 4-2.

Nerita micheloti

*Loriol, P. de. 1867. In P. de Loriol and E. Pellat.
Monographie paleontologique et geologique de l'etages
portlandien du department de l'Yonne. Bull. Soc. Sci.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

N. micheloti, cont.
Hist. Nat. l'Yvonne 21, p. 35.

Nerita minima

*Credner, H. 1864. Die Pteroceras-Schichten (Aporrhais-Schichten) der Umgebung von Hannover. Z. Deutschen Geol. Gesell- 16, p. 223.

Nerita minnesotensis

*Bergquist, H. R. 1944. Cretaceous of the Mesabi Iron Range, Minnesota. J. Paleontol. 18, p. 9, 25. [Colerine Formation, Cretaceous, Calumet, Minnesota]

Nerita minor

*Sandberger, F. 1863. Die Conchylien des Mainzer Teriat-beckens, p. 158.

Nerita minutaeformis

*Schauroth, K. F. v. 1865. Verzeichniss der Versteine-rungen im Herzogl. Naturalienkabinet zu Coburg, p. 133.

Nerita monodon

*Ferussac, A. E. 1823. In G. P. Deshayes In A. E. Ferussac. H. N. g. et p. Moll., wrapper of livr. 20. [n. et f.]

Nerita montegrottensis

*Gregorio, A. 1907. Microfaune des Thermes Euganees (Abano, Bottaglia, etc.). Ann. Geol. Paleontol. 32, p. 9.

Nerita morellii

*Bellardi, L., and G. Michelotti. 1841. Mem. Roy. Acc. Sci. Torino 3, p. 165.

Nerita morio

*Dujardin, G. 1837. Mem. Soc. Geol. France 2, p. 280. [invalid name, see N. glibertii]

Nerita mosae

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 7.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita moulinsii

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 40.

Nerita multicarinata

*Doncieux, L. 1908. Catalogue descriptif des fossiles nummulitiques de l'Aude et de l'Hérault. Ann. Univ. Lyon, n.s., 22, p. 235.

Nerita munita

*Forbes, E. 1856. Trans. Geol. Soc. London, Ser. 2, 7, p. 122.

Nerita munsteri

*Hornes, M. 1855. Über die Gastropoden und Acephalen der Hallstätter Schichten. Denksch. Kaiserlichen Akad. Wissenshc. 9., p. 40. [Tertiary Europe]

Nerita muricata

*Young, ., and . Bird. 1828. Geol. Surv. Yorksh. Coast, ed. 2, p. 249.

Nerita narbonensis

*Noulet, J.-B. 1858. De l'âge géologique de la formation lacustre de Narbonne et de Sigean (Aude). Mem. Acad. Imp. Sci., Inscript. Belles-lettres Toulouse, Ser. 5, 2, p. 421.

Nerita naticiformis

*Piette, E. 1855. Observations sur les étages inférieurs du terrain jurassique dans les départements des Ardennes et de l'Aisne. Bull. Soc. Geol. France, Ser. 2, 12, p. 1110.

Nerita nebrodensis

*Gemmellaro, G. G. 1866. Naticidae e Neritidae del terreno giurassico del nord di Sicilia. G. Sci. Nat. Econom. 2, p. 178.

Nerita nodosa

*Stephenson, L. W. 1947. Prof. Pap. U.S. Geol. Surv. 210E, p. 180. [Upper Cretaceous, USA]

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita nodoso-costata

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 192.
[Natica from Index Animalium 2]

Nerita novem sulcata

*Molina, G. I. 1810. Sag. Stor. Nat. Chili, ed. 2, p. 177.

Nerita nucleus

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 301.
[Neritina? from Index Animalium 2]

Nerita nuda

*Piette, E. 1855. Observations sur les etages inferieurs du terrain jurassique dans les departements de Ardennes et de l'Ainse. Bull. Soc. Geol. France, Ser. 2, 12, p. 1110.

Nerita nudispira

*Piette, E. 1855. Observations sur les etages inferieurs du terrain jurassique dans les departements de Ardennes et de l'Ainse. Bull. Soc. Geol. France, Ser. 2, 12, p. 1110.

Nerita oligopleura

*Dall, W. H., and W. H. Ochsner. 1928. Tertiary and Pleistocene Mollusca from the Galapagos Islands. Proc. California Acad. Sci. 17, p. 114.

Nerita orbignyana

*Cotteau, G. 1854. Etudes sur les mollusques fossiles du departement de l'Yonne. Premier fascicule comprenant l'introduction et le prodrome, p. 30.

Nerita ornata

*Forbes, E. 1856. Report on the fossil invertebrata from southern India, collected by Mr. Kaye and Mr. Cunliffe. Trans. Geol. Soc. London, Ser. 2, 7, p. 121.

Nerita ornata

*Pcelincev, V. 1927. Mem. Com. Geol. Leningrad, n.s., 172, p. 203. [Jurassic Caucasus]

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita ornatissima

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 192.

Nerita ovata

*Roemer, F. A. 1836. Nerst. Norddeutsch. Oolith. 3,
p. 156.

Nerita oviformis

*Forbes, E. 1856. Report on the fossil invertebrata from
southern India, collected by Mr. Kaye and Mr. Cunliffe.
Trans. Geol. Soc. London, Ser. 2, 7, p. 122.

Nerita oyula

*Buvignier, A. 1843. Mem. Soc. Philom. Verdun, 2, p. 241.

Nerita pagoda

Whitfield, R. P. 1891. Observations on some Cretaceous
fossils from the Beirut district of Syria, in the collection
of the American Museum of Natural History, with descriptions
of some new species. Bull. American Mus. Nat. Hist. 3,
p. 432. [Abeih, Syria]

Nerita palaeochroma

*Buvignier, A. 1843. Mem. Soc. Philom. Verdun, 2, p. 241.

Nerita palaeomorpha

*Gemmellaro, G. G. 1891. La fauna dei Calcari con Fusulina
della valle del Fiume Sosio nella Provincia di Palermo. G.
Sci. Palerm. 19, p. 77. [Fusulina Limestone, Province of
Palermo]

Nerita palauensis

Ladd, H. S. 1966. Chitons and gastropods (Haliotidae
through Adeorbidae) from the western Pacific islands. Geol.
Surv. Prof. Pap. 531, p. 56. [Palau Limestone, Auluptagel
Is., Palau, late Miocene]

Nerita pallidula

*Risso, A. 1826. H. N. Europe 4, p. 151.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita papilionacea

*Valenciennes, A. 1832. In Humbolt. Voy. Inter. Amea.
(Obs. Zool. II, p. 264).

Nerita paretii

*Gemmellaro, G. G. 1866. Naticidae e Neritidae del terreno giurassico del nord di Sicilia. G. Sci. Nat. Econom. 2, p. 183.

Nerita pecosensis

*Stanton, T. W. 1947. Studies of some Comanche pelecypods and gastropods. Geol. Surv. Prof. Pap. 211, p. 211.
[Comanche, Lower Cretaceous]

Nerita pentastoma

*Deshayes, G. P. 1864. Description des animaux sans vertebres decouverts dans le bassin de Paris...3, p. 17.

Nerita petersi

*Gemmellaro, G. G. 1869. Sulla fauna calcario a Terebratula janitor del nord di Sicilia. G. Sci. Nat. Econom. 4, p. 149.

Nerita picta

*Ferussac, A. E. 1823. In G. P. Deshayes In A. E. Ferussac. H. N. g. et p. Moll., wrapper of livr. 20.
[n. et f.]

Nerita pisca

*Gemmellaro, G. G. 1891. La fauna dei Calcari con Fusulina della valle de Fiume sosio nella Provinicia di Palermo. G. Sci. Palerm. 19, p. 77. [Fusulina Limestone, Province of Palermo]

Nerita plebeia

*Reuss, A. E. 1846. Verst. boehm. Kreidef 2, p. 112.

Nerita plexa multistriata

*Abrard, R. 1942. Arch. Mus. Paris, Ser. 6, 18, p. 55.
[Pleistocene, French Somaliland]

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita plicatula

*Anton, H. E. 1839. Verzeichniss der Conchylien..., p. 30.

Nerita platonis

*Daudin, F. M. 1803. Sonnini's Buffon Rept. 4, p. 324.

Nerita poliana

*Chiaje, S. d. 1830. Mem. s. vert. Napoli, Tab. pro V & VI, pl. 104.

Nerita ponderosa

*Piette, E. 1855. Observations sur les etages inferieurs du terrain jurassique dans les departements de Ardennes et le ll'Aisne. Bull. Soc. Geol. France, Ser. 2, 12, p. 1094.

Nerita ponderosa

*Cossmann, M. 1912. Rectification de nomenclature. Rev. Crit. Paleozool. Paris 16, p. 213. [N. ponderosa de Lar. 1890 non Piette, 1855 = N. cercicinensis Cossm., 1912]

Nerita prevosti

*Gemmellaro, G. G. 1866. Naticidae e Neritidae del terreno giurassico del nord di Sicilia. G. Sci. Nat. Econom. 2, p. 180.

Nerita prevostiana

*Partsch, . 1828. In C. Pfeiffer. Naturg. deutsch. land-moll. 3, p. 49.

Nerita pricei

*Shumard, B. F. 1858. In B. F. Shumard and G. C. Swallow. Descriptions of new fossils from the coal measures of Missouri and Kansas. Trans. Acad. Sci. St. Louis 1, p. 202.

Nerita prinzingeri

*Hornes, M. 1855. Über einige Gastopoden aus den östlichen Alpen. Denksch. Kaiserlichen Akad. Wissenschaft. 10, pl. 175. [Tertiary]

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita prototypa

*Phillips, J. 1848. Mem. Geol. Surv. Gt. Britian 2,
p. 358.

Nerita pseudo-concava

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 7.

Nerita pseudo-costata

d'Orbigny, A. 1849. Prodrome de paleontologie...1, p. 264.

Nerita pulchella

*Buvignier, A. 1852. Statistique geologique, mineralo-
gique, minerallurgique et paleontologique du departement de
la Meusse, atlas, p. 30.

Nerita pulla

*Roemer, F. A. 1836. Verst. Norddeutsch. Oolith. 3,
p. 155.

Nerita punctata

*Piette, E. 1855. Observations sur les etages inferieurs
du terrain jurassique dans les departements des Ardennes et
de l'Aisne. Bull. Soc. Geol. France, Ser. 2, 12, p. 1110.

Nerita punctata

*Smith, W. D. 1906. Preliminary geological reconnaissance
of the Laboo Mountains, Batangas Province. Philippine J.
Sci. 1, p. 628.

Nerita pustulata

*Gemmellaro, G. G. 1966. Naticidae e Neritidae del terreno
giuassico del nord del Sicilia. G. Sci. Nat. Econom. 2,
p. 183.

Nerita quadrangulata

*Weaver, D. W., and R. M. Kleinpell. 1963. Systematic
catalogue and register of localities. In Oligocene
biostratigraphy of the Santa Barbara Embayment, California.
Publ. Dep. Geol. Univ. California 43, p. 183. [Sacate-
Gaviota formation, Oligocene, Los Olivas Quadrangle,
California]

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita rapula

*Nardo, G. D. 1847. Sin. mod. spec. Lag. Veneto, p. 97.

Nerita rhenana

*Thomas, C. 1845. Jarb. Ver. Nat. Nassau 2, p. 161.

Nerita ringicula

*Philippi, R. A. 1851. Centuria quarta testaceorum novorum. Z. Malakozool. 8, p. 86.

Nerita rugosa

*Hoeninghaus, F. W. 1830. Jarhb. Min., p. 467. [nomen nudum]

Nerita rugosa

*Anton, H. E. 1839. Verzeichniss der Conchylien...., p. 29.

Nerita rugosa

*Morris, J., and J. Lycett. 1854. A monograph of the Mollusca of the Great Oolite, chiefly from Minchinghamton and the coast of Yorkshire, part 1, p. 56. [Jurassic fossil]

Nerita rustica

*Nardo, G. D. 1847. Sin. mod. spec. Lag. Veneto, p. 97.

Nerita sadebekki

*Schmidt, M. 1905. Über Oberen Jura im Pommern. Beiträge zur Stratigraphie und Palaontologie. Abh. Geol. Landesanst., N.F., 41, p. 183.

Nerita sancti-stephani

*Cossmann, M., and . Peyrot. 1919. Bordeaux Actes Soc. Linn. 70, p. 40. [Neogene France]

Nerita savii

*Gemmellaro, G. G. 1869. Sulla fauna calcario a Terebratula janitor del nord di Sicilia. G. Sci. Nat. Econom. 4, p. 148.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita saxoides

*Nardo, G. D. 1847. Sin. mod. spec. Lag. Veneto, p. 97.

Nerita scalaris

*Seeley, H. 1861. Notes on Cambridge palaeontology. Ann. Mag. Nat. Hist., Ser. 3, 7, p. 288.

Nerita scilla

*Nardo, G. D. 1847. Sin. mod. spec. Lag. Veneto, p. 95.

Nerita semilugubris

*Deshayes, G. P. 1864. Descriptions des animaux sans vertebres decouvertes dans le bassin de Paris...3, p. 15.

Nerita semi-luna

*Piette, E. 1856. Notice sur les gres d'Aiglemont et de Rimogne. Bull. Soc. Geol. France, Ser. 2, 13, p. 205.

Nerita semi-luna

*Terquem, O., and E. Piette. 1865. Le lias inferieur de l'est de la France comprenant la Meurthe, la Moselle, le Grand-duche de Luxembourg, la Belgique et la Meuse. Mem. Soc. Geol. France, Ser. 2, 8, p. 45.

Nerita semipleura

*Twenhofel, W. H. 1924. The geology and invertebrate palaeontology of the Comanchean and "Dakota" formations of Kansas. Bull. State Geol. Surv. Kansas 9, p. 56.

Nerita semistriata

*Meunster, G. v. 1840. Beitr. Petref. 3, p. 83.

Nerita semisulcata

*Brocchi, . 1821. Bibl. Ital. 24, p. 225.

Nerita semisulcata

*Gemmellaro, G. G. 1866. Naticidae e Neritidae del terreno giurassico del nord di Sicilia. G. Sci. Nat. Econom. 2, p. 182.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita sena

*Contraine, F. 1835. Bull. Acad. Roy. Brux. 2, p. 390.

Nerita sigaretina

*Buvignier, A. 1843. Mem. Soc. Philom. Verdun 2, p. 241.

Nerita sinuosa

*Sowerby, J. DeC. 1821. The mineral conchology of Great Britain;...3, p. 32.

Nerita spadæ

*Gemmellaro, G. G. 1896. Sulla fauna calcario a Terebratula janitor del nord di Sicilia. G. Sci. Nat. Econom. 4, p. 149.

Nerita spirata

*Sowerby, J. DeC. 1825. The mineral conchology of Great Britain;...5, p. 93.

Nerita stefaninii

*Thalem, . 1937. Palaeontogr. Ital. Pisa 37, p. 9.
[Cretaceous Syria]

Nerita stragulata

*Megerle, J. K. 1828. In C. Pfeiffer. Naturg. deutsch. Land-Moll. 3, p. 49

Nerita stremmei

*Dietrich, W. O. 1914. Die Gastropoden der Tandaguru-schichten, der Aptstufe und der Oberkreide im sudlichen Deutsch-Ostafrika. Berlin Arch. Biontol. 3, p. 126.
[Jurassic, East Africa]

Nerita striata

*Defrance, . 1825. Dict. Sci. Nat. 34, p. 478.

Nerita striata

*Fleming, J. 1828. Hist. Brit. Anim., p. 319.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita striata

*d'Orbigny, A. 1842. In Sogea, Cuba (Moll. II), p. 50.

Nerita subantonii

*Aliev, G. A. 1958. New Gastropoda from Lower Cretaceous deposits of Little Caucasus (Azerbaijan). Izv. Akad. Nauk Azerbaidzh. Geol. Geog. 2, p. 36. [in Russian]

Nerita subcarinata

*Matheron, P. 1843. Catal. meth. foss. B. du Rhone, p. 229; Rep. Trav. Soc. Stat. Marseille 4, p. 301.

Nerita subcaronis

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 41.

Nerita subcompacta

*Trechmann, C. T. 1927. The Cretaceous shales of Jamaica. Geol. Mag. 64, p. 35.

Nerita subconcava

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 40.

Nerita subcornea

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 40.

Nerita subcostata

*Goldfuss, . 1832. In Dechen In De la Beche. Handb. Geogr., p. 523.

Nerita subgranulosa

*Nagao, T. 1928. Palaeogene fossils of the island of Kyushu, Japan. Sci. Rept. Tohoku Univ. 9, p. 117.

Nerita subintermedia

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 40.

Nerita sublineolata

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 346.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

Nerita subornata

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 301.

Nerita subpicta

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 39.

Nerita subpisiformis

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 40.

Nerita subuplicata

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 40.

Nerita sub-savii

*Pcelincev, V. 1927. Mem. Com. Geol. Leningrad, n.s., 172, p. 251.

Nerita subsulcosa

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 39.

Nerita subvirginea

d'Orbigny, A. 1852. Prodrome de paleontologie...3, p. 40.

Nerita sulcatina

*Buvignier, A. 1852. Statistique geologique, mineralogique, minéralurgique et paleontologique du département de la Meuse, atlas, p. 30.

Nerita sulcatina

*Gemmellaro, G. G. 1869. Sulla fauna calcario a Terebratula janitor del nord di Sicilia. G. Sci. Nat. Econom. 4, p. 147.

Nerita sanurensis

*Oppenheimer, P. 1906. Zur Kenntnis des ältertiaret Faunen in Ägypten. 2. Lieferung... Palaeontographica 30, p. 226. [Tertiary Egypt]

Nerita tampaensis

*Dall, W. H. 1916. A contribution to the invertebrate

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]

N. tampaensis, cont.

fauna of the Oligocene beds of Flint River, Georgia. Proc. U.S. Nat. Mus. 51, p. 487.

Nerita teinostoma

*Ryckholt, P. de. 1860. Melanges paleontologiques. Mem. Couronnes Mem. Savants Etrangers, publies par l'Academie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique 1, plate 25.

Nerita theodorii

*Kuhn, O. 1936. Die Fauna des Amaltheentons (Lias 8) in Franken. N. Jarhb. Min. Geol. Palaont. Stuttgart 75, p. 287.

Nerita thersites

*Bayan, F. 1870. Sur les terrains tertiaires de la Venetie. Bull. Soc. Geol. France, Ser. 2, 27, p. 457.

Nerita thurmanni

*Loriol, P. de, and E. Koby. 1890. Etudes sur les Mollusques des Couches Coraligenes inferieures du Jura Bernois. Part I. Abh. Schw. Pal. Ges. 16, p. 102.

Nerita triangulata oregonensis

*Merriam, C. W., and F. E. Turner. 1937. The Capay middle Eocene of northern California. Univ. California Publ. Geol. 24(6), p. 104.

Nerita tricarinata

*Lamarck, . 1804. Ann. Mus. H. N. (Paris), 5, p. 94.

Nerita tricarinata

*Pcelincev, V. 1927. The Jurassic and Lower Cretaceous fauna of the Crimea and Caucasus. Mem. Com. Geol. Lenin-grad, n.s., 172, p. 152.

Nerita tuberosissima

*Montagu, G. 1808. Test. Brist. Suppl., p. 150.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]Nerita turbinata

*Sharpe, D. 1850. Q. J. Geol. Soc. London 6, p. 192.

Nerita umzambiensis

*Woods, . 1906. Ann. S. Af. Mus. 4, p. 311 [Cretaceous Ponodland]

Nerita undata pseudoplica

*Strausz, L. 1960. Neue Namem und neue Formen unter den Miozanmollusken. Foldt. Kozl. 90, p. 348, 352.

Nerita veldiensis

*Roemer, F. A. 1839. Verst. Norddeutsch. Oolth. Nachtrag, p. 47.

Nerita venusta

*Muenster, G. v. 1840. Beitr. zur Petref. 3, p. 84. [Triassic]

Nerita vexillum

*Megerle, J. K. 1824. Ges. Nat. Fr. Berlin, Verh. I(4), p. 221.

Nerita vicina

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 301.

Nerita vokesi

*Durham, J. W. 1944. Univ. California Publ. Geol. 27(5), p. 118 [Oligocene north-western Washington]

Nerita wabrensis

*Buvignier, A. 1852. Statistique geologique, mineralogique, minéralurgique et paleontologique du département de la Meuse, atlas p. 30.

Nerita washingtoniana

*Weaver, C. E., and K. van W. Palmer. 1922. Fauna from the Eocene of Washington. Univ. Washington Publ. Geol. 1(3), p. 28.

TABLE 2. (Continued) LIST OF FOSSIL NERITA SPECIES
[* = not seen]Nerita wrightiana

*Cotteau, B. 1854. Etudes sur les mollusques fossiles du departement de l'Yonne. Premier fascicule comprenant l'introduction et le prodrome, p. 30.

Nerita zebra

*Bruguiere, J. G. 1792. Actes Soc. H. N. Paris 1, p. 126.

Nerita zigzag

*Michelotti, G. 1840. Ann. Sci. Lomb.-Ven. 10, p. 144.

Nerita zonaria

d'Orbigny, A. 1850. Prodrome de paleontologie...2, p. 312.
[now Neritina]

Nerita zumoffeni

*Delpay, G. 1940. Les gasteropodes mesozoiques de la Region libanaise. Notes Mem. Haut-Comm. Republ. Franc. Syrie et Au Liban 3, p. 41.

TABLE 3

NERITA SPECIES NAMES DESCRIBED MORE THAN ONCE
 [+ = fossil; n.n. = nomen nudum]

<u>aculeata</u>	Muller, 1774 Gmelin, 1791
<u>angulosa</u>	Meuschen, 1787 (n.n.) Recluz, 1842
<u>antiquata</u>	Recluz, 1841 Recluz, 1853
<u>australis</u>	Gmelin, 1791 Wood, 1828 (n.n.)
<u>cardinalis</u>	Bolten, 1798 (n.n.) LeGuillou, 1841
<u>chrysostoma</u>	Karsten, 1789 Bolten, 1798 (n.n.) Recluz, 1841
<u>cornea</u>	Linnaeus, 1758 Gratélop, 1828
<u>corona</u>	Linnaeus, 1758 deBlainville, 1824
<u>deformis</u>	Sowerby, 1840 +Gabb, 1864
<u>dubia</u>	Gmelin, 1791 Turton, 1932
<u>eburnea</u>	Bolten, 1798 (n.n.) Hoeninghaus, 1831
<u>funiculata</u>	Menke, 1850 Reeve, 1855
<u>grisea</u>	Meuschen, 1787 (n.n.) Reeve, 1855
<u>grossa</u>	Linnaeus, 1758 +Bronn, 1848
<u>guttata</u>	Meuschen, 1787 (n.n.) Recluz, 1841

TABLE 3. (Continued) NERITA SPECIES NAMES DESCRIBED MORE THAN ONCE. [+ = fossil; n.n. = nomen nudum]

<u>lineata</u>	Muller, 1774 Meuschen, 1787 (n.n.) Gmelin, 1791
<u>listeri</u>	Pfeiffer, 1840 Recluz, 1841
<u>littoralis</u>	Linnaeus, 1758 Risso, 1826
<u>maculata</u>	Salis, 1793 Bolten, 1798 (n.n.) Pease, 1867 Doncieux, 1908
<u>marmorata</u>	Meuschen, 1787 (n.n.) +Link, 1807 Reeve, 1855
<u>minima</u>	LeGuillou, 1841 +Credner, 1864
<u>minuta</u>	Muller, 1774 Sowerby, 1825
<u>olivacea</u>	Bolten, 1798 (n.n.) Recluz, 1842
<u>ornata</u>	Sowerby, 1823 +Forbes, 1856 Pcelincev, 1927
<u>patula</u>	Wood, 1828 (n.n.) Recluz, 1841
<u>pica</u>	Meuschen, 1787 (n.n.) Gould, 1859 Lishcke, 1869
<u>plumbea</u>	Meuschen, 1787 (n.n.) Wood, 1828 (n.n.)
<u>pulchella</u>	Wood, 1828 (n.n.) Recluz, 1843 +Buvignier, 1852

TABLE 3. (Continued) NERITA SPECIES NAMES DESCRIBED MORE THAN ONCE. [+ = fossil; n.n. = nomen nudum]

<u>punctata</u>	...	Muller, 1774 Meuschen, 1787 (n.n.) Dillwyn, 1817 Quoy and Gaimard, 1834 +Piette, 1855 +Smith, 1906
<u>reticulata</u>	...	Meuschen, 1787 (n.n., p. 272) Meuschen, 1787 (n.n., p. 274) Karsten, 1789 Quoy and Gaimard, 1834
<u>rugosa</u>	...	Gmelin, 1791 +Hoeninghaus, 1830 +Anton, 1839 +Morris and Lycett, 1854
<u>striata</u>	...	Burrow, 1815 +Defrance, 1825 +Fleming, 1828 +d'Orbigny, 1842
<u>sulcatina</u>	...	+Buvignier, 1852 +Gemmellaro, 1869
<u>versicolor</u>	...	Meuschen, 1787 (n.n.) Gmelin, 1791
<u>violacea</u>	...	Meuschen, 1787 (n.n.) Gmelin, 1791
<u>viridis</u>	...	Linnaeus, 1758 Meuschen, 1787 (n.n.)

LITERATURE CITED

- Council of Biology Editors. 1972. CBE style manual, third edition. American Institute of Biological Sciences, Washington, D. C. 297 p.
- Gmelin, J. F. 1791. *Systema naturae*, 13th edition. G. E. Beer, Leipzig.
- Hanley, S. 1855. *Ipsa Linnaei conchylia. The shells of Linnaeus, determined from his manuscripts and collections.* Williams and Norgate, London.
- Hemming, F. 1958. Official index of rejected and invalid works in zoological nomenclature. First installment: Names 1-58. International Trust for Zoological Nomenclature, London. 14 p.
- International Commission for Zoological Nomenclature. 1985. International code of zoological nomenclature, third edition. International Trust for Zoological Nomenclature, London. 338 p.
- Linnaeus, C. 1758. *Systema naturae*, 10th edition. Laurentii Salvii, Stockholm.
- Ruhoff, F. A. 1980. Index to the species of Mollusca introduced from 1850 to 1870. Smithsonian Contrib. Zool. 294:1-640.
- Sherborn, C. D. 1902-1933. *Index animalium sive index nominum quae ab A.D. MDCCCLVIII generibus et speciebus animalium imposita sunt.* London.